

The air quality perceptions of the residents of Bayview, Mossel Bay

Johann P. Schoeman¹ and De Wet Schutte²

¹28 Maroela Street, Mossel Bay, Western Cape, 6500, South Africa, jschoeman@edendm.co.za

²Cape Peninsula University of Technology, Post Office Box 652, Cape Town 8000, South Africa

Abstract

Background: In developing countries, it often occurs that little attention is given to air pollution emissions due to a lack of proper town planning, household combustion processes, energy production and the continuous growth in the transport sector (Norman *et al.*, 2007:783). There is an increase in urban air pollution in most of the major cities of developing countries which is amplified by population growth and industrialization (World Resource Institute, 1998, 1999:1). Air pollution studies are not complete, and may fail if the quality of life and the perceptions of the studied community are not taken into consideration. This paper investigates the air quality perceptions of a high income residency surrounded by industrial activities and Mossel Bay was rated as to have potentially poor air quality by the South African Department of Environmental Affairs and Tourism.

Methods: A cross-sectional survey was carried out in Bayview, Mossel Bay. The perceptions of the respondents were collected by a structured questionnaire. Components of perceptions that were tested included general opinion regarding air quality, visual perceptions of air quality, type of pollutants such as smoke and dust, perceptions regarding the source of air pollution, perceptions regarding the municipal health institution controlling air quality in Bayview, etc. These perceptions were investigated by age, gender, socio-economic status etc.

Conclusion: The findings of the study indicated that various factors, such as visual impacts, type of pollutants, role of the municipal health institution governing air quality, influence the air perceptions of the Bayview residents.

Keywords

air pollution, perceptions, air quality, monitoring, visual impacts, exposure, social status, local knowledge

Introduction

In developing countries it often occur that little attention is given to air pollution emissions due to a lack of proper town planning, household combustion processes, energy production and the continuous growth in the transport sector (Norman *et al.*, 2007). Most of the major cities of developing countries show an increase in urban air pollution which is amplified by population growth and industrialisation (World Resource Institute, 1998, 1999).

One of the major air quality problems in South Africa is poor regional and town planning, often driven by political ideologies where residential areas are developed adjacent or next to industrial areas (Matookane *et al.*, 2004). Mossel Bay was no exception to this rule and Bayview, one of the suburbs of this developing town is surrounded by industrial activities. The Mossel Bay region was also classified by the Department of Environmental Affairs as to have potentially poor air quality. One of the reasons for this classification was the petro-chemical industry situated 10km west of the Bayview residency. This industry triggered other industrial activities in the region. Considerable emphasis is placed on the monitoring of air pollutants that pose a risk and important aspects such as personal and community opinions on air quality and the effect on their health are often overlooked (Hunter *et al.*, 2003).

The development of perceptions about environmental pollution is a complex process due to the culmination of a wide spectrum of possible sensory inputs that could converge into perceptions (Bickerstaff, 2003). In this respect it is clear that attitudes and concerns that an individual might have concerning the environment can play an important role in what is perceived as an environmental hazard (Stenlund *et al.*, 2009). Therefore, a general definition of risk perception is that it includes a person's attitude, behaviour, beliefs, opinions and concerns for the environmental hazard (Stenlund *et al.*, 2009). Bickerstaff (2003) takes this notion further as he argues that risk perceptions include the wider cultural and social disposition of people towards dangers and the disadvantages thereof. From this approach it is clear that perceptions regarding air pollution are predominantly based on the visual and chemo-sensory indicators, and showed as being useful indicators for possible air pollution when one has to mediate between environmental exposure and health (Stenlund *et al.*, 2009).

Hunter *et al.*, (2003) recorded that public stress related to the potential health effects of air pollution on individuals and their families increased over the last decades. Hunter *et al.*, (2003) furthermore mentioned various studies have been published on air pollution and the prevalence of respiratory diseases, the link between these two variables are still not clear, simply because of the lack of scientifically acceptable control over the

variables in a real world setting. In this respect it was found that local knowledge could play an important role in shaping the people’s perceptions of environmental risks (Scammel *et al.*, 2009). Research on perceptions of environmental risks showed that local context and experience plays an important role in the definition and perceptions of environmental risks (Bickerstaff, 2004). Added to this, it is also known that risk perceptions are also influenced by an individual’s perception of the credibility of the environmental health institution that has to take care of his/her health (Scammel *et al.*, 2009).

Hyslop (2009) argued that the human physiology and psychology is particularly sensitive for visual inputs and that the observable, impacts to a large extent dictate how people perceive the world around them. Although man usually tends not to pay attention to his environment, it still consciously or subconsciously, influences how he sees life (Hyslop, 2009). However, since the 1960’s there been a growing sensitivity for air pollution, mainly as a result of increased publicity, with the result of increased awareness regarding air quality (Hyslop, 2009).

This paper investigates the perceptions regarding the air quality of Bayview, middle to upper income residential suburb of Mossel Bay in South Africa. Being surrounded by industrial activities, Mossel Bay was rated by the South African Department of Environmental Affairs as having potentially poor air quality.

Methods

Bayview is middle to upper income residential suburb of Mossel Bay and is almost surrounded by industrial activities that include petrochemical storage tanks, a coal burning food industry, light industrial activities and also railway activities operating steam and diesel locomotives. A systematic sample of 114 plots (total 483) was selected from a cadastral map of Bayview. The sample size was calculated by means of Statcalc statistical software using a confidence interval of 5 points and a confidence level of 95% accuracy. A Politz frame was used to randomly select a respondent among the members of the selected household that qualifies for this study. Respondents that qualified for selection by means of the Politz frame had to be 18 years of age and older, usually sleep in the household and lived in the house since January 2009. The normal ethical conditions like anonymity, informed consent and voluntary participation applied during data collection. The research study and questionnaire was approved by the Cape Peninsula University of Technology Ethical Committee and Applied Sciences Higher degrees Committee.

The target population for this study consisted of all the residents that were 18 years and older and that have resided in Bayview for at least one year prior to the data collection for this research project. A questionnaire was designed to yield information regarding the perceptions of the residents regarding the air quality of the area. The questionnaire was based on literature research making use of the dendrogram-technique which is a conceptual framework highlighting the research questions on the lowest level of the dendrogram. The questionnaire

included sections on quality of life and perceptions. The section on perceptions included questions such as length of stay in Bayview, visibility of pollution and industrial activity, evaluating the general air quality of bayview, perception on industrial activity with biggest impact, perception on pollution control at industrial level and authority level, etc. Risk perceptions are also influenced by a person’s perception of the worthiness of the health institution that has to protect that person’s health (Scammel *et al.*, 2009). In order to test this theory the question of how the respondent perceived the health institution responsible for the health of the Bayview residents, was posed.

Independent variables in this evaluation were the respondent’s age, gender, and period of residency, self reported health status, health consciousness, and exposure to polluted air in the workplace. Dependent variables in this study were their general perceptions of the air quality and visibility of air pollutants in Bayview. The results of the questionnaire were coded and analysed with the StatSoft Statistica version 9 software package. Chi square analysis was used to search for statistical significant relationships between independent and dependent variables.

Results and discussion

Table 1 shows the sample demographics for the 114 respondents.

Table 1: The age distribution of respondents

Age	n	%
18 - 23	5	4.4
24 - 30	3	2.6
31 - 45	20	17.5
46 - 60	41	36
61 +	45	39.5
Total	114	100.0

The age distribution indicated that the age group of 61+ was the highest age group in Bayview. Only 4.4% respondents were between 18 and 23 years old. The age profile thus indicated a high age profile.

Table 2: The comparison between male and female of the respondents

Sex	n	%
Male	52	45.6
Female	62	54.3
Total	114	99.9

Females were the predominant sex with a sex distribution of 54.3% female and 45.6% male respondents.

Table 3: The length of residency in Bayview

Length of residency in Bayview	n	%
0 - 5 years	32	28.1
6 - 10 years	19	16.7
11 - 20 years	44	38.6
21 years and longer	19	16.6
Total	114	100.00

Thirty-eight point six percent of the population resided in Bayview between 11-20 years and 16.6% for longer than 21 years. Based on this information it was derived that the population would have known the residency well and that their local knowledge would have been set.

Monitoring results

The study also took the air quality status of Bayview into account by studying the air quality data of the Department of Environmental Affairs and Development Planning ambient air quality station located in Bayview. The results indicated no exceedences of the South African National Ambient Air Quality Standards for the criteria pollutants. The annual average concentrations of the measured criteria pollutants can be observed in figure 1.




Figure 1: The annual average concentrations of NO₂, PM₁₀, O₃, C₆H₆ and SO₂ for the Bayview residency from 1st October 2008 till 30 September 2009

The data of the DEADP monitoring station were captured and compared to the 90% completeness against the USEPA and SANS1929 standards. See table 4.

Table 4: The annual average of completeness of the captured data for the period 1 October 2009 to 30 September 2009 (South Africa, DEADP, 2009:10)

Pollutant	SO ₂	NO ₂	O ₃	PM ₁₀
% collected	96.8%	98.9%	92.6%	97.6%

Perceptions

Almost two thirds 63.1% (n=72) of the respondents indicated that they perceived that the air quality was good, with a further 13.2% (n=15) who rated the air quality as excellent, whilst 19.3% (n=22) thought it was average (Table 5). The minority of respondents rated the air quality as poor to very poor (4.3%, n= 5).

Table 5: The perceptions of the Bayview respondents on the status of air quality

Opinion of the Bayview air quality	n	%
Excellent	15	13.2
Good	72	63.1
Average	22	19.3
Poor	4	3.5
Very poor	1	0.8
Total	114	99.9

Results of the logistic regression showed that the following factors may influence perceptions of air quality among the residents of Bayview, Mossel Bay. Although various influences

such as age, gender, length of stay, etc were tested, only the factors with statistical significance (P<0.05) are mentioned in this paper.

Visual impacts

Because visual impact plays an important role in the formation of perceptions regarding air quality, respondents were asked what they consider to be the main contributors to air pollution in Bayview. The results show that the perceived main contributor to pollution in the area is the Food factory (34.2%, n=39), followed by (19.3%, n=22) that perceive it is one of the local transport companies, followed by Petrochemical storage facility and the railway loco, both on (15.8%, n=18). (See Table 6)

Table 6: Perceptions of Bayview respondents regarding the surrounding industries with the biggest air quality impact

Industries with the major impact on Bayview air quality	n	%
Not answered	3	2.6
Petrochemical Storage facility	18	15.8
Local transport company	22	19.3
Railway loco	18	15.8
Tank farms	8	7.0
Oil tankers	4	3.5
Food Factory	39	34.2
Seal island	2	1.8
Total	114	100

Exposure to air pollution

The study found a statistically significant correlation (p=0.035) between the perception of the air quality of Bayview and the perception of the influence of the industries on the respondents' health (Table 7). It was found was that the respondents (80%, n=12) that perceived the air quality as excellent also indicated that the surrounding industries do not have a negative impact on their health. The same tendency was found with the respondents that perceived the air quality as good, when 61.1% (n=44) indicated that the surrounding industries do not have a negative influence on their health. This tendency was further confirmed when the majority (54.5%, n=12) among the 22 respondents (19.3%) that rated the air quality as average, indicated that they perceived that the industries do have a negative impact on their health.

Table 7: The correlation between the perceptions of air quality and the perceptions of the impact of the industries on the health of the respondents

Perceptions of air quality	Industries do influence my health		Industries does not influence my health		Unsure, if industries influence my health		Total	
	n	%	n	%	n	%	n	%
Excellent	1	6.7	12	80.0	2	13.3	15	100.0
Good	15	20.8	44	61.1	13	18.1	72	100.0
Average	12	54.5	6	27.3	4	18.2	22	100.0
Poor	3	75.0	0	0.0	1	25.0	4	100.0
Very poor	1	100.0	0	0.0	0	0.0	1	100.0
Total	32	28.1	62	54.4	20	17.5	114	100.0

Pearson Chi-square: 22.8375, df=8, p=.003582

Social status

Perceptions are also influenced by the environment, availability of information and socio-economic characteristics (MackKerron et al., 2009). Previous studies indicated that the more wealthy communities have the perception that air quality imposes a bigger threat to the less fortunate communities (Bickerstaff, 2004). The social status of the Bayview suburb was high. It was derived from their income level, work level and educational level. Thirty seven point eight percent, (n=43) respondents were professional, 39.4%, (n=45) had an annual income of R180 001 and higher and 43.8%, (n=50) had a degree or diploma. This research found a correlation between qualification and perceptions.

Table 8: The correlation between perceptions of air quality and the industrial emission control

Qualification	Yes, industries emission control		No, industries emission control		Unsure, industries emission control		Total	
	n	%	n	%	n	%	n	%
< Grade 12	10	55.6	2	11.1	11.1	33.3	18	100.0
Grade 12	13	28.3	13	28.3	28.3	43.5	46	100.1
> Grade 12	8	16.0	17	34.0	34.0	50.0	50	100.0
Total	31	27.2	32	28.0	51	44.8	114	100.0

Pearson Chi-square: 10.9681, df=4, p=.026931

Eighteen of the respondents (15.7%) n=18 had qualifications lower than grade 12. Of those, 11.1% n=2 indicated that the industries don't do enough to control emissions, 33.3% n=6 was unsure and 55.6% n=10 indicated adequate emission control. Forty six (40.4%) respondents had grade 12. Of those, 28.3%, (n=13) indicated that industries don't do adequate emission control and almost half (43.5%, n=20) indicated that they don't know. Forty three point eight percent n=50 had a qualification higher than grade 12 (diploma or degree) of which 34%, (n=17) indicated that industries do not control emissions adequately and 50%, (n=25) indicated that they don't know. The perceptions of emission control fluctuated over educational qualification. The higher the qualification thus the higher the perception that industries do not implement adequate emission control. These groups were also unsure if the industries are doing emission control.

Risk perceptions and the perceptions of the air quality authority regulating air quality in Bayview

It was clear that the majority of the respondents that classified the air quality as excellent and good (n=15 and 72 respectively) did not know how to judge the health authority (46.7% and 36.1% respectively) (Table 9). Of these two groups, the second highest percentage (26.7%, n=4 and 34.7%, n=25) respectively also judged the health institution as good.

Table 9: The correlation between the perceptions of the Bayview respondents and the perception of the air quality authority

Perception air quality	Health institutions excellent		Health institutions good		Health institutions average		Health institutions poor		Health institutions don't know		Total	
	n	%	n	%	n	%	n	%	n	%	n	%
Excellent	3	20.0	4	26.7	1	6.7	0	0.0	7	46.7	15	100.1
Good	1	1.4	25	34.7	18	25.0	2	2.8	26	36.1	72	100.0
Average	0	0.0	5	22.7	6	27.3	4	18.2	7	31.8	22	100.0
Poor	0	0.0	1	25.0	0	0.0	2	50.0	1	25.0	4	100.0
Very poor	0	0.0	0	0.0	0	0.0	1	100.0	0	0.0	1	100.0
TOTAL	4	3.5	35	30.7	25	22.0	9	7.9	41	36.0	114	100.1

The influence of visibility on risk perceptions

On the question if any of the industries were visible from the respondents dwelling, 59.6%, (n=68) answered yes and 40.4%, (n=46) answered no. The majority respondents could see an industry from their home (Table 10).

Table 10: The visibility of the industries from the respondent's dwellings

Visibility of the industries	n	%
Yes	68	59.6
No	46	40.4
Total	114	100.00

There were 53.5%, (n=61) respondents that could see smoke from their dwellings (Table 11). More than two thirds 70.1%, n=80 of respondents experienced or saw dust from their dwellings.

Thirteen point two percent respondents (n=15) that perceived the air quality as excellent, the minority or 46.7%, (n=7) experienced or saw smoke from their houses. Almost two thirds or (63.2%, n=72) respondents that perceived the air quality to be good, the minority or 45.8%, n=33 could see smoke from their dwellings. Four respondents perceived the air quality as poor and all of them, n=4 could see smoke from their dwellings.

Table 11: The correlation between perceptions and the experiencing or visibility of smoke.

Perception of air quality	See smoke		Don't see smoke		Total	
	n	%	n	%	n	%
Excellent	7	46.7	8	53.3	15	100.0
Good	33	45.8	39	54.2	72	100.0
Average	17	77.3	5	22.7	22	100.0
Poor	4	100.0	0	0.0	4	100.0
Very poor	0	0.0	1	100.0	1	100.0
Total	61	53.5	53	46.5	114	100.0

Pearson Chi-square: 11.6079, df=4, p=.020523

Conclusion

This study found the higher the social status in terms of qualification, thus the lower the perception regarding the risk. Smoke posed a significant impact on the perceptions of the respondents. The importance of visibility was also emphasized as the industry that was most visible was also perceived as the industry with the biggest threat or impact. There is a correlation between general air quality perceptions and industrial air quality impact. There is a correlation between air quality perceptions and the perceptions of the air quality authority regulating air quality of that specific population.

References

Bickerstaff, K. 2004. *Risk Perception research: socio-cultural perspectives on the public experience of air pollution*. 2004. Environment International volume 30 (2004). Science direct.

www.elsevier.com/locate/envint Pages 827-840. [11 August 2009].

Carter, S., Williams, M., Paterson, J., Lusitini, L. 2008. *Do perceptions of neighbourhood problems contribute to maternal health? Findings from the Pacific Islands Families study.* (PIF) Study, AUT University, Auckland, New Zealand. Health & Place. 2008. www.elsevier.com/locate/healthplace Pages. 622-630. [15 May 2009].

Encyclopedia Britannica. 2009. Encyclopedia Britannica Online. 2009. <http://www.search.eb.com/eb/article-9110465> Pages. 1-3. [22 May 2009].

Hunter, P., Davies, M., Hill, K., Whittaker, M., Sufi, F. 2003. The prevalence of self-reported symptoms of respiratory disease and community belief about the severity of pollution from various sources. September 2003. *The International Journal of Environmental Health Research* 13(3). Taylor and Francis Health Sciences. London UK. Pages 227-338.

Hyslop, N. 2009. *Impaired visibility: the air pollution people see.* *Atmospheric Environment* 43 (2009). Science Direct. www.elsevier.com/locate/atmosenv Pages 182-185. [22 May 2009].

MacKerron, G., Mourato, S. 2009. *Life satisfaction and air quality in London.* 2009. *Ecological Economics* Vol. 68 (2009) Science Direct. www.elsevier.com/locate/ecocon Pages 1441-1450. [20 July 2009].

Matooane M., John, J., Oosthuizen, R en Binedell, M. 2004. Vulnerability of South African Communities to Air Pollution. Proceedings of the 2004 Conference of the South African Institute of Environmental Health, Durban, 22-27 February 2004.

Norman, R., Eugene Cairncross, Jongikhaya Witi, Debbie Bradshaw. 2007. *Estimating the burden of disease attributable to urban outdoor air pollution in South Africa in 2000.* *South African Medical Journal*, 97(7) August 2007. Pages 782-790.

Scammel, M., Senier, L., Darrah-Okike, J., Brown, P. 2009. Tangible evidence, trust and power: *Public perceptions of community environmental health studies.* *Social Science & Medicine*, 68(2009). Science Direct. www.elsevier.com/locate/socscimed Pages 143-146. [22 May 2009].

Stenlund, T., Lide, E., Andersson, K., Garvill, J., Nordin, S. 2009. *Annoyance and health symptoms and their influencing factors: A population-based air pollution intervention study.* *Public Health*. 2009. Science Direct. www.elsevierhealth.com/journals/pubh Pages 339-345. [15 May 2009].

World Resource Institute. Health and Environment, 1998, 1999. Health Effects of Air Pollution. <http://www.wri.org/wr-98-99/airpoll.htm> [11 May 2009]