

Selecting an appropriate method for expressing S locus F-box-S2 recombinant protein

Jahanshah Ashkani and D.J.G. Rees

Abstract:

A single locus (*S* locus) including at least two linked genes (female and male determinants) genetically controls the gametophytic self-incompatibility (GSI) in apple, which has evolved to avoid self-fertilization. There has been extensive work done on the female determinant of self-incompatibility, which has led to the determination of the tertiary structure of *S*-RNase. However, the tertiary structure of male determinant (*S* locus F-box, SLF/SFB) remains unresolved, which could mainly be due to difficulties associated with its expression in the recombinant expression systems. In addressing this, we have evaluated several *in vivo* (prokaryotic and eukaryotic) and *in vitro* expression systems for their efficiency in the expression of apple SLF2. The most successful expression of SLF2 (1 mg/ml) was achieved in *E. coli* using the synthesized gene in a high salt culture and applying heat shock before induction of culture. We therefore present an approach for the efficient expression of *S* locus F-box recombinant proteins for future functional and structural studies.

1. Introduction

The growing economic importance of fruit crops and the problems associated with their commercial production has placed the molecular basis of self-incompatibility (SI) under much attention. There has been extensive work done on the female determinant of self-incompatibility, which has led to the determination of the tertiary structure of S3-RNase in *Pyrus pyrifolia* [1,2]. However, the male determinant (*S* locus F-box, SLF/SFB) structure remains unresolved, which could mostly be due to difficulties related to its expression in the recombinant expression systems.

Although, gene expression is a strong tool for studying biological processes, choosing an appropriate expression system based on the characteristics and deliberated application of a recombinant protein is critical to obtain adequate quantities of the protein in a timely manner [3,4]. Many expression systems are available but no single system has yet been developed that works for all proteins [5–11].

Escherichia coli is a popular organism in *in vivo* studies due to its relatively easy and inexpensive genetic manipulation, scalability of the experiments, low cost, fast growth, high-density cultivation and its ability to express labeled proteins [5,9,12]. However, certain limitations may be experienced primarily at the initiation of translation [13–17].

To overcome such limitations certain *E. coli* strains can overexpress rare tRNAs (for AGG, AGA, CUA, AUA, CCC, and GGA codons) and hence significantly improve gene expression [18–20]. Furthermore, genes are designed in certain cases to match host bias or to maximize Codon Adaptation Index (CAI) and contribute towards an improved expression [13,21]. In addition, synthetic gene sequences designed by codon engineering can create an optimal heterologous gene expression and are considered important in strategically improve protein expression [22,23].

Besides *E. coli*, several other recombinant protein expression systems are available in eukaryotes, including fungal, mammalian, amphibian, insect, plants and yeast cells. Among the available host expression systems, *Escherichia coli* and cultivated mammalian cells are considered most widely used [24–26]. Advantages of using a mammalian expression system include its ability for post-translational modifications (PTMs) such as correct protein folding and glycosylation. The protein yield of mammalian expression systems can be optimized by the use of right cell line, expression vector, promoter elements and transfection efficiency of the mammalian cells [25–27].

Another commonly used and well-established technique in recombinant protein expression is the Transient Expression Arabidopsis Mesophyll Protoplasts (TEAMP), which is appropriate for studying various physiological processes. The TEAMP is a highly useful and sensitive system because it does not need a large number of protoplasts, sterile techniques or complex culture medium [28]. Expression methods have become good alternatives for cell-based protein production. Firstly, it is easily adapted for high-throughput and automated procedures. Secondly, the availability of various prokaryotic and eukaryotic cell lysates provides a flexible choice for high-level protein expression, especially those that are difficult to express under an *in vivo* system. Finally, cell-free systems are capable of synthesizing large protein–protein complex mixtures in a single reaction, which makes them an exploitable system for developing proteomic tools [29–32]. Cell-free protein synthesis has many applications in the fields of genomics, proteomics, and synthetic biology while allowing for increased productivity and various post-translational modifications [33,34].

Despite great progress in understanding the complex non-self recognition between male and female determinants, a comprehensive structural study is still necessary to understand the biochemical basis of differential interactions between male and female determinants (*i.e.* SLF/SFB and SRNase) in *Rosaceae*. Here, we aim to evaluate the capability and applicability of several prokaryotic and eukaryotic expression systems in expressing the *S* locus F-box-S2 recombinant protein. Since F-box proteins represent a big family of proteins in plants (*e.g.*, the *Arabidopsis thaliana* genome encodes more than 700 F-box proteins) and multiple F-box genes with more than 60% similarity have been recognized at the *S*-loci of the families that own S-RNase-based SI [35–37], the information provided here not only improves our understanding of SRNase-based self-incompatibility, but also lays the foundation for the functional and structural studying of the interactions between F-box proteins and their substrates.

2. Material and methods

2.1 Recombinant protein expression in *E. coli* and purification

For the purpose of construct preparation for recombinant protein expression; nucleic acid extraction, PCR amplification, and DNA sequencing were done (Supplementary information 1). The SLF2-pGEX-6P-2, SLF2-pET21a and SLF2-pETM-41 constructs were prepared by digestion of SLF2-pGEM-T Easy constructs using double digestion with BamHI and XhoI, and NcoI and BamHI respectively, where the restriction enzyme sites were added to SLF2 codon sequence (Accession Number: DQ422811) using designed forward and reverse primers (Table S1) by PCR technique. The restriction enzyme assays were used to check the availability of SLF2 in the constructs. Separate conditions were considered with combining vector plasmids, cell lines and IPTG concentrations (0.1 mM–1.0 mM), while the temperature was decreased to (25–30 °C) following IPTG induction and cells were harvested between 4 and 24 h with 4 h intervals. The absence of SLF2 expression was further assessed using blotting techniques (Supplementary information 2.1). Gene synthesis was carried out by GenScript (<http://www.genscript.com>) after a wide variety of factor optimization using OptimumGene™ algorithm to generate a single sequence gene that can reach the highest possible level of expression in *E. coli* (Supplementary information 2.2, Fig. S1 and S2). The localization of the GST-SLF2opt in *E. coli* was analyzed by using osmotic shock procedure and an anionic detergent (Sarkosyl) and 8 M Urea. GST-SLF2opt was expressed in *E. coli* C41 (DE3) pLysS while grown in LB containing 0.5 M NaCl and induced with 0.1 mM IPTG. Furthermore, solubility and localization of GST-SLF2opt were assessed (Supplementary information 2.3).

2.2 Recombinant protein expression in mammalian cells

The SLF2-FX-6xHis-pcDNA3.1/Zeo(+) was constructed by digestion of SLF2-pGEM-T Easy constructs using double digestion of NheI and NotI restriction enzymes added to SLF2 through the use of designed forward and reverse primers (Table S1) using the PCR technique. SLF2 expression was conducted using pcDNA3.1/Zeo(+) expression vector as a 6xHis fusion protein with a designed Factor Xa cleavage site before the 6xHis tag at the C-terminal end. CHO cells were grown in complete Hams F12 at 37 °C in a humidified atmosphere containing 5% CO₂. Secondary cultures were used for transfection. Transfected cells were grown in normal growth media after growing in selective media containing 500–600 µg/ml zeocin for a week. The cells were lysed mechanically using RIPA buffer by passing the pellet through a syringe needle on ice. To detect expression of SLF2 recombinant fusion protein into CHO cells, Western blot analysis was performed while an anti-His antibody was used to specifically bind to 6xHis tag in C-terminal of the SLF2 on PVDF Western blotting membrane (Supplementary information 3).

2.2 Recombinant protein expression in *Arabidopsis thaliana* protoplasts

The SLF2-FX-6xHis-pCAMBIA1-305.1 was constructed by the digestion of SLF2-pGEM-T-Easy constructs using double digestion of NdeI and BamHI restriction enzymes whose sites were added to SLF2 (Table S1) using PCR technique. SLF2 expression was conducted using pCAMBIA1305.1 expression vector as a 6xHis fusion protein with a designed Factor Xa cleavage site before the 6xHis tag at the C-terminal end (Table S1). The PEG-mediated method was used to transfect a large amount of SLF2-FX-6xHis-

pCAMBIA1305.1 plasmid DNA (30–40 µg) into the healthy and successful isolated protoplasts. Subsequently, the samples were loaded on the SDS-PAGE and the Western blot technique was applied to confirm the expression of SLF2 in the protoplasts (Supplementary information 4).

2.3 Cell-free (in vitro) recombinant protein expression

The pIVEX (*In Vitro* EXpression) vectors are specially designed for gene expression *in vitro*. These vectors contain an ampicillin resistance gene, T7 promoter, and 6xHis-tag that facilitate the high level of expression, detection and purification of proteins. The pIVEX2.3d (3560 bp) contains an uncleavable 6xHis-tag at the C-terminal end. The pIVEX2.4d (3583 bp) contains a 6xHis-tag at the N-terminal end, which is cleavable with Factor Xa. The SLF2-pIVEX2.3d and SLF2-pIVEX2.4d were constructed by the cloning of SLF2 gene isolated from apple *var.* Golden Delicious into pIVEX2.3d using NcoI and SmaI and pIVEX2.4d expression vector using NotI and SmaI restriction sites added to the SLF2 codon sequence (Table S1) using PCR technique, respectively. To find the best condition for maximum expression per reaction both constructs of SLF2-pIVEX2.3d and SLF2-pIVEX2.4d were used as templates in concentrations of 5 µg/ml, 8.5 µg/ml and 10 µg/ml along with 0.27% (v/v) of cell extract and standard reaction mix. Samples were incubated at temperatures of 25, 30 and 37 °C in an Eppendorf Thermomixer R for 4–12 h while shaking at 250 rpm. For the protein expression assessment, samples were centrifuged at 10,000 × *g* for 15 min and 50 µl of suspension was loaded on the SDS-PAGE and analyzed by Western blotting (Supplementary information 5). When the incubation time increased from 4 to 12 h, the expression of the SLF2 gene was increased accordingly (data not shown).


Results

3.1. Recombinant protein expression in *E. coli*

The main focus of this study has been the over-expression of *S* locus F-box-S2 pollen factor of apple *var.* Golden Delicious (SLF2) in a bacterial expression system to provide sufficient amount of SLF2 for protein studies. To achieve this goal, several cloning vectors (pGEX-6P-2, pET21a and pETM-41) were tested in different bacterial cell lines (BL21(DE3)-pLysS, C41(DE3)-pLysS, and C43(DE3)-pLysS). However, no significant expression for SLF2 was found.


3.2. Recombinant protein expression in mammalian cells

SLF2 expression was conducted using pcDNA3.1/Zeo(+) expression vector as a 6xHis fusion protein with a designed Factor Xa cleavage site before the 6xHis tag at the C-terminal end (Table S1). CHO cells were grown in complete Hams F12 at 37 °C in a humidified atmosphere containing 5% CO₂. The results of this study proved successful although only very faint expression of this gene was observed in the CHO cells lysates (Fig. 1).




(a) Untransfected CHOs.

(b) Untransfected CHOs under the selective conditions.



(c) Transfected CHOs under the selective conditions.



(d) Western blot analysis of the proteins extracted from the CHO cells.

Fig. 1. CHO Transfection with SLF2-FX-6xHis-pcDNA3.1/Zeo (+) and protein expression in CHO cells. a) Positive control sample, Un-transfected CHOs. b) Negative control sample with addition of selective media (containing 600 µg/ml Zeocin). c) The growing number of successfully transfected cells after the addition of selective media (containing 600 µg/ml Zeocin). d) Western blot analysis of SLF2 expression in CHO cells using Anti-His antibody on PVDF membrane.

3.3 Recombinant protein expression in mesophyll protoplasts of *Arabidopsis*

We further examined *Arabidopsis* protoplasts as an alternative expression host for expressing the plant-derived SLF2 protein. Despite the successful use of protoplast expression system by other scientists, the results indicate a very faint signal on the PVDF membrane at 45 kDa (Fig. 2) while no detectable band was observed on the SDS-PAGE suggesting a very low level of expression of the SLF2 protein.

3.4 Cell free (*in vitro*) recombinant protein expression

To find the best condition for maximum expression per reaction both constructs of SLF2-pIVEX2.3d and SLF2-pIVEX2.4d were used as templates in several DNA concentrations (5–10 µg/ml) under temperatures between 25 and 37 °C for 4–12 h. The result of the optimization shows that the concentration of 8.5 µg/ml DNA in reaction while incubated at 30 °C for 12 h, yields the best results for the expression of SLF2 *in vitro* (Fig. 3). Nonetheless, the *in vitro* expression system did not meet our priorities due to its shortcomings such as it being costly and its low protein yield (~ 25–30 µg/ml).


3.5 Recombinant protein expression using synthetic SLF2 gene (SLF2opt) in *E. coli*

We carried out SLF2 gene synthesis through GenScript ([http:// www.genscript.com](http://www.genscript.com)) after a broad range of factors were optimized using OptimumGene™ algorithm to generate a single sequence gene that can reach the highest possible level of expression in *E. coli*. The SLF2opt-pGEX-6P-2 was constructed using BamHI and XhoI restriction enzyme sites and the construct was transformed into the C41 (DE)pLysS cells. The growth of bacterial cultures containing SLF2opt-pGEX- 6P-2 was measured at OD600 every 30 min after the IPTG induction and the result showed that the GST-SLF2opt


protein inhibits the growth of *E. coli* cells (data not shown) and an over-expression of GST-SLF2opt fusion protein (~70 kDa) was shown in the insoluble fraction (Fig. 4). To enhance the solubility of GST-SLF2opt fusion protein, growth medium (LB) was replaced with a high salt growth medium (LB containing 0.5 M NaCl) and a heat-shock step (incubating the culture in a 47 °C water-bath for 30 min) was applied before IPTG induction when the OD600 of the culture was between 0.8-1. Sixteen different parameters containing IPTG concentrations (0.1, 0.25, 0.5 and 1 mM) and overnight (16 h) incubation temperatures (18, 20, 25 and 30 °C) were examined using transformed BL21, BL21-CodonPlus, C41 (DE3)-pLysS and C43 (DE3)-pLysS cell lines. The Dot blot immunoassay was used as a technique for rapid testing of the parameters for maximum expression of GST-SLF2opt fusion. The result of the Dot blot assay shows a significant signal for C41 (DE3)-pLysS and C43 (DE3)-pLysS cell lines when compared to BL21 and BL21-CodonPlus. Furthermore, the most significant signal belongs to C41 (DE3)-pLysS cells that were induced with 0.1 mM IPTG concentration and further incubated at 20 °C overnight (Fig. 5).

3.5.1. Cellular localization of GST-SLF2opt fusion protein

The localization of the GST-SLF2opt in *E. coli* was assessed using osmotic shock, anionic detergent (Sarkosyl) and 8 M Urea. GST-SLF2opt was expressed in *E. coli* C41 (DE3)-pLysS while grown in LB containing 0.5 M NaCl and induced with 0.1 mM IPTG. This resulted in the yield of 1 mg/l of soluble protein (proteins in periplasmic, inner membrane and cytoplasmic fractions) measured at A260. However, the biggest amount of expressed proteins was presented in the form of inclusion bodies (Fig. 6). We further used a GST column for affinity purification of the GST-SLF2opt recombinant protein. To release the GST domain from GST-SLF2opt, PreScission protease was used in a dialysis bag while stirring in 1 liter cleavage buffer at 4 °C overnight. Following the cleavage, about three-quarter of SLF2opt proteins was precipitated in the absence of the fusion tag in the dialysis bag (data not shown) and the amount of soluble SLF2opt was decreased to 0.25 mg/l.



(a) Isolated Arabidopsis mesophyll protoplasts.



(b) Western blot analysis of the proteins extracted from the protoplasts.

Fig. 2. Recombinant protein expression of SLF2 in Arabidopsis protoplasts. a) Protoplasts isolation from *Arabidopsis thaliana* cell suspension and b) Western blot analysis of the proteins extracted from the protoplasts.




Fig. 3. Western blot result of *in vitro* SLF2 recombinant protein expression. First lane shows protein marker and lanes 2-4 show 5 µg/ml, 8.5 µg/ml and 10 µg/ml DNA concentration at 25 °C, lanes 5-7 show 5 µg/ml, 8.5 µg/ml and 10 µg/ml DNA concentration at 30 °C and lanes 8-10 show 5 µg/ml, 8.5 µg/ml and 10 µg/ml DNA concentration at 37 °C, respectively.

Fig. 7 illustrates the elution fraction containing GST-SLF2opt fusion protein (~70 kDa) obtained from the protein purification step before and after 3C protease cleavage. Two bands in Lane C correspond to GST (26.6 kDa) and the SLF2opt (43 kDa) proteins, which is indicative of the complete cleavage of GST-SLF2opt fusion protein.


We used Anion exchange chromatography with a pre-packed MonoQ 10/100 GL column on AKTA Chromatography System (GE Healthcare Life Sciences) to separate GST proteins from SLF2opt recombinant protein in the cleaved sample. The chromatogram shows a number of small peaks containing SLF2opt recombinant protein and a big peak containing GST proteins as confirmed on an SDS-PAGE (Fig. 8).

Discussion

It is postulated that the lack or slow progress in structural aspects of the *Rosaceae* self-incompatibility may be due to the existing limitations on the expression of SLF/SFB in the heterologous expression systems. Accordingly, we evaluated several expression systems for their efficiency in expressing adequate quantity of the *S* locus-F-box-S2 (SLF2) recombinant protein. Most of these methods had limitations regarding their yield, cost, and time efficiency. However, the successful expression of SLF2 was achieved in *E. coli* using the synthesized gene. Furthermore, SLF2 was expressed in a soluble form using high salt culture and heat-shock application before the induction of culture.

Here we report that the production of an SLF2 protein optimized for expression in an *E. coli*-based system is shown to be the most desirable technique for high yield of SLF2. Using a high-salt media along with heat shock regimes before induction significantly contributed to the production of more soluble recombinant proteins even though a large proportion of GST-SLF2opt fusion protein was still in the inclusion bodies. This necessitates future studies to focus on the optimization of the conditions that give the maximum amount of GST-SLF2opt in a soluble form. Additionally, more work needs to be carried out to optimize the purification of SLF2, as we encountered protein precipitation during the purification steps.

In a study on *Antirrhinum* (family: *Plantaginaceae*) Qiao et al. showed that the expression of AhSLF-S2 was only made possible following the separation of the protein into 6xHis-AhSLF-S2-N and 6xHis-AhSLF-S2-C for the pull-down assay [38]. However, it is well established that the tertiary structure of a protein is essential in understanding its function [39]. We, therefore, aimed at expressing SLF2 as a complete protein molecule, however despite the use of several *E. coli* expression systems, no significant expression was observed. This lack of expression in *E. coli* could be due to un-optimized factors responsible for the efficiency of heterologous protein production such as biased codon usage and mRNA secondary structure [12,40]. As a result, mammalian cell expression systems were investigated as transient expression systems that allow extra-chromosomal amplification of plasmids and hence permit more plasmid copies to persist in the transfected cells during the production phase for a significant increase in gene expression. The large-scale expression of recombinant proteins in mammalian cells has been shown to generate sufficient amounts of recombinant protein for early production and high-throughput screening in functional studies [41].



(a) SDS-PAGE of the total bacterial lysates. (b) Western blot analysis of the total bacterial lysates.

Fig. 4. Solubility analysis of GST-SLF2opt fusion protein. Soluble (liquid fraction after centrifuging, lane B) and in-soluble (pellet dissolved in 8 M urea, lane C) fractions of total bacterial lysates using (a) SDS-PAGE and (b) Western blot analysis.

However, the expression of SLF/SFB recombinant proteins in mammalian cells requires optimization of the technique to improve the transfection procedure through increasing speed and reducing the costs, all of which can possibly make mammalian cells more suitable for SLF/ SFB recombinant protein expression. We further assessed the use of protoplasts isolated from plants as means for the transient expression of recombinant proteins. Protoplasts retain their cell identity and show high transformation efficiency while requiring low maintenance [42]. Despite the successful use of protoplast expression system as a recombinant expression system by other scientists such as Wu et al., very faint expression of the SLF2 protein was observed in this study.

Another system examined in this study is the cell-free/*in vitro* expression system that can synthesize proteins in a high speed and accurate manner. The results obtain from this analysis suggest very low levels of expression. The *in vitro* system, by eliminating the need to manipulate whole cells in culture is a suitable technique for automated, high-throughput expression of a large number of proteins. However, the very low productivity *i.e.* less than 30 µg protein/ml reaction of this system limits its applicability and make it unfeasible compared to cell- based systems.

Subsequently, the optimization of the *E. coli* system to express SLF/ SFB gene was considered using artificial gene synthesis. Considering that each organism carries its own bias in the usage of the 61 available codons [15] in some cases expression is limited primarily to the initiation of translation [13,14,43,44]. In addition to codon bias, mRNA secondary structures are identified which affect the expression of the encoded proteins [15]. A synthetic gene sequence by codon engineering is an important strategy for enabling optimal heterologous gene expression through making DNA templates ready to be expressed in any heterologous host [22,23]. This method verified to significantly contribute towards the expression of SLF/SFB in *E. coli*. As such it may be deduced that among the expression systems under study artificial gene synthesis optimized for over-expression in *E. coli* provides the best technique for the optimal expression of SLF/SFB and may further be used in future structural studies while folding and activity of the expressed protein must be evaluated. Having said that, the protocols provided for the mammalian cells, Arabidopsis protoplast and *in vitro* expression systems might also be used for studies with low protein yield requirement such as functional studies using co-immunoprecipitation and single-molecule pull-down assays.

Here, we have reported on the efficacy of a broad range of expression systems in expressing SLF2 recombinant protein of apple *var.* Golden Delicious, with artificial gene synthesis proving to be the best approach. The findings of our study can guide future structural-based studies with the foremost goal of better understanding the mechanism of SI in *Rosaceae*.

		IPTG concentration (mM)																			
		0.1				0.25				0.5				1							
Temperature (°C)	18																				
	20																				
	25																				
	30																				
		BL21				BL21-CodonPlus				C41 (DE3)-pLysS				C43 (DE3)-pLysS							

Fig. 5. Detection of suitable parameters for GST-SLF2opt expression in *E. coli* cell lines using Dot blot. Sixty-four different parameters including IPTG concentrations (0.1–1 mM), overnight incubation temperatures after induction (18–30 °C) and different *E. coli* cell lines (BL21, BL21-CodonPlus, C41 (DE3)-pLysS and C43 (DE3)-pLysS) were analyzed on PVDF Western blotting membrane using anti-GST antibody.




Fig. 6. Cellular localization of GST-SLF2opt fusion protein. First lane represents marker and lanes A-D illustrate the amount of GST-SLF2opt in the inclusion bodies and outer membrane (A), periplasmic (B), inner membrane (C) and cytoplasmic (D) fractions in *E. coli*.






Fig. 7. PreScission[™] protease cleavage of the GST-SLF2opt fusion protein. Lane A shows the molecular weight marker, while lanes B and C show the un-cleaved and cleaved GST-SLF2opt fusion proteins, respectively. Two bands in Lane C are corresponded to the GST (26.6 kDa) and the SLF2opt (43 kDa) proteins.



(a) Chromatogram showing anion exchange chromatography of the SLF2opt.



(b) SDS-PAGE of anion exchange fractions for SLF2opt and GST.

Fig. 8. Anion exchange purification and SDS-PAGE of anion exchange fractions for SLF2opt and GST. **a)** The chromatogram shows two peaks; a smaller wide peak, corresponding to SLF2opt recombinant protein and a sharp peak, corresponding to the GST proteins. **b)** Lane A shows protein marker, lane B shows GSTs in GST domain peak and lanes C-G show SLF2opt in different fraction of SLF2opt peak.

Conflict of interest declaration

None.

Financial disclosure

Not available.

Acknowledgements

This project was supported by National Research Foundation (www.nrf.ac.za) and Agriculture Research Council (www.arc.agric.za) of South Africa.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.btre.2017.06.005>.

References

- [1] T. Matsuura, M. Unno, H. Sakai, T. Tsukihara, S. Norioka, Purification and crystallization of Japanese pear S-RNase associated with gametophytic self-incompatibility, *Acta Crystallogr. D Biol. Crystallogr.* 57 (1) (2001) 172–173.
- [2] T. Matsuura, H. Sakai, M. Unno, K. Ida, M. Sato, F. Sakiyama, S. Norioka, Crystal structure at 1.5 Å resolution of *Pyrus pyrifolia* pistil ribonuclease responsible for gametophytic self-incompatibility, *J. Biol. Chem.* 276 (48) (2001) 45261–45269.
- [3] A. Vilcinskas, C. Becker-Pauly, W. Stöcker, Insect cells for heterologous production of recombinant proteins, *Insect Biotechnology* vol. 2, Springer Netherlands, 2011, pp. 197–209 (Chapter 11).
- [4] W.H. Brondyk, Selecting an appropriate method for expressing a recombinant protein, in: R.B. Richard, P.D. Murray (Eds.), *Methods in Enzymology*, vol. 463, Academic Press, 2009, pp. 131–147.
- [5] P. Balbás, A. Lorence, *Recombinant Gene Expression: Reviews and Protocols* vol. 267, Springer Science & Business Media, 2004.
- [6] A. Villaverde, M. Mar Carrio, Protein aggregation in recombinant bacteria: biological role of inclusion bodies, *Biotechnol. Lett.* 25 (2003) 1385–1395.
- [7] G. Georgiou, P. Valax, Expression of correctly folded proteins in *Escherichia coli*, *Curr. Opin. Biotechnol.* 7 (2) (1996) 190–197.
- [8] A. Middelberg, Preparative protein folding, *Trends Biotechnol.* 20 (2002) 437.
- [9] H.P. Sorensen, K.K. Mortensen, Advanced genetic strategies for recombinant protein expression in *Escherichia coli*, *J. Biotechnol.* 115 (2) (2005) 113–128.
- [10] F.W. Studier, A.H. Rosenberg, J.J. Dunn, J.W. Dubendorff, Use of T7 RNA Polymerase to direct expression of cloned genes, *Methods Enzymol.* 185 (1990) 60–89.
- [11] G.D. Davis, C. Elisee, D.M. Newham, R.G. Harrison, New fusion protein systems designed to give soluble expression in *Escherichia coli*, *Biotechnol. Bioeng.* 65 (4) (1999) 382–388.
- [12] N.H. Tolia, L. Joshua-Tor, Strategies for protein coexpression in *Escherichia coli*, *Nat. Methods* 3 (1) (2006) 55–64.
- [13] C. Gustafsson, S. Govindarajan, J. Minshull, Codon bias and heterologous protein expression, *Trends Biotechnol.* 22 (7) (2004) 346–353.
- [14] G. Kudla, A.W. Murray, D. Tollervey, J.B. Plotkin, Coding-sequence determinants of gene expression in *Escherichia coli*, *Science* 324 (5924) (2009) 255–258.
- [15] D. Lorimer, A. Raymond, J. Walchli, M. Mixon, A. Barrow, E. Wallace, R. Grice, Burgin, L. Stewart, Gene Composer: database software for protein construct design, codon engineering, and gene synthesis, *BMC Biotechnol.* 9 (1) (2009) 36.
- [16] J.T. Wilson, L.B. Wilson, V.B. Reddy, C. Cavallero, P.K. Ghosh, J.K. deRiel, G. Forget, S.M. Weissman, Nucleotide sequence of the coding portion of human alpha globin messenger RNA, *J. Biol. Chem.* 255 (7) (1980) 2807–2815.
- [17] G. Modiano, G. Battistuzzi, A.G. Motulsky, Nonrandom patterns of codon usage and of nucleotide substitutions in human alpha- and beta-globin genes: an evolutionary strategy reducing the rate of mutations with drastic effects? *Proc. Natl. Acad. Sci. U. S. A.* 78 (2) (1981) 1110–1114.
- [18] B.J. Del Tito Jr., J.M. Ward, J. Hodgson, C.J. Gershter, H. Edwards, L.A. Wysocki, F.A. Watson, G. Sathe, J.F. Kane, Effects of a minor isoleucyl tRNA on heterologous protein translation in *Escherichia coli*, *J. Bacteriol.* 177 (24) (1995) 7086–7091.

- [19] J.F. Kane, Effects of rare codon clusters on high-level expression of heterologous proteins in *Escherichia coli*, *Curr. Opin. Biotechnol.* 6 (5) (1995) 494–500.
- [20] N.A. Burgess-Brown, S. Sharma, F. Sobott, C. Loenarz, U. Oppermann, O. Gileadi, Codon optimization can improve expression of human genes in *Escherichia coli*: A multi-gene study, *Protein Expr. Purif.* 59 (1) (2008) 94–102.
- [21] P.M. Sharp, W.-H. Li, The codon adaptation index—a measure of directional synonymous codon usage bias, and its potential applications, *Nucleic Acids Res.* 15 (3) (1987) 1281–1295.
- [22] L. Stewart, A.B. Burgin, Whole gene synthesis: a gene-o-matic future, *Front. Drug Des. Discov.* 1 (1) (2005).
- [23] S.J. Maerkl, Next generation microfluidic platforms for high-throughput protein biochemistry, *Curr. Opin. Biotechnol.* 22 (2010) 1–7.
- [24] P.A. Grimsrud, D.L. Swaney, C.D. Wenger, N.A. Beauchene, J.J. Coon, Phosphoproteomics for the masses, *ACS Chem. Biol.* 1 (2010) 105–119.
- [25] F. Krammer, J. Pontiller, C. Tauer, D. Palmberger, A. Maccani, M. Baumann, R. Grabherr, Evaluation of the Influenza A replicon for transient expression of recombinant proteins in mammalian cells, *PLoS One* 5 (10) (2010) e13265.
- [26] J. Pontiller, A. Maccani, M. Baumann, I. Klancnik, W. Ernst, Identification of CHO endogenous gene regulatory elements, *Mol. Biotechnol.* 45 (3) (2010) 235–240.
- [27] L. Baldi, D. Hacker, M. Adam, F. Wurm, Recombinant protein production by large-scale transient gene expression in mammalian cells: state of the art and future perspectives, *Biotechnol. Lett.* 29 (5) (2007) 677–684.
- [28] S.-D. Yoo, Y.-H. Cho, J. Sheen, Arabidopsis mesophyll protoplasts: a versatile cell system for transient gene expression analysis, *Nat. Protoc.* 2 (7) (2007) 1565–1572.
- [29] F.-H. Wu, S.-C. Shen, L.-Y. Lee, S.-H. Lee, M.-T. Chan, C.-S. Lin, Tape-Arabidopsis Sandwich—a simpler Arabidopsis protoplast isolation method, *Plant Methods* 5 (1) (2009) 16.
- [30] E.C. Cocking, A Method for the isolation of plant protoplasts and vacuoles, *Nature* 187 (4741) (1960) 962–963.
- [31] M. He, Cell-free protein synthesis: applications in proteomics and biotechnology, *New Biotechnol.* 25 (2–3) (2008) 126–132.
- [32] M. He, In vitro protein expression: an emerging alternative to cell-based approaches, *New Biotechnol.* 28 (3) (2011) 209–210.
- [33] M.W. Nirenberg, J.H. Matthaei, The dependence of cell-free protein synthesis in *E. coli* upon naturally occurring or synthetic polyribonucleotides, *Proc. Natl. Acad. Sci. U. S. A.* 47 (10) (1961) 1588–1602.
- [34] T.A. Nguyen, S.S. Lieu, G. Chang, An *Escherichia coli*-based cell-free system for large-scale production of functional mammalian membrane proteins suitable for X-ray crystallography, *J. Mol. Microbiol. Biotechnol.* 18 (2) (2010) 85–91.
- [35] J.M. Gagne, B.P. Downes, S.H. Shiu, A.M. Durski, R.D. Vierstra, The F-box subunit of the SCF E3 complex is encoded by a diverse superfamily of genes in Arabidopsis, *Proc. Natl. Acad. Sci. U. S. A.* 99 (17) (2002) 11519–11524.
- [36] E.P. Risseuw, T.E. Daskalchuk, T.W. Banks, E. Liu, J. Cotelesage, H. Hellmann, M. Estelle, D.E. Somers, W.L. Crosby, Protein interaction analysis of SCF ubiquitin E3 ligase subunits from Arabidopsis, *Plant J.* 34 (6) (2003) 753–767.
- [37] Z. Hua, X. Meng, T.-h. Kao, SLF-like proteins reveals its unique function in S-RNase?based self-incompatibility, *Plant Cell* 19 (11) (2007) 3593–3609.

- [38] H. Qiao, F. Wang, L. Zhao, J. Zhou, Z. Lai, Y. Zhang, T.P. Robbins, Y. Xue, The F-box protein AhSLF-S2 controls the pollen function of S-RNase-based self-incompatibility, *Plant Cell* 16 (9) (2004) 2307–2322.
- [39] M.F. Perutz, Species adaptation in a protein molecular biosystemseccule, *Mol. Biol. Evol.* 1 (1) (1983) 1–28.
- [40] E.L. Rashdy, M. Redwan, The optimal gene sequence for optimal protein expression in *Escherichia coli*: principle requirements, *Arab J. Biotechnol.* 9 (3) (2006) 493–510.
- [41] R. Kunaparaju, M. Liao, Sunstrom N-A: Epi-CHO, an episomal expression system for recombinant protein production in CHO cells, *Biotechnol. Bioeng.* 91 (6) (2005) 670–677.
- [42] J. Sheen, Signal transduction in maize and Arabidopsis mesophyll protoplasts, *Plant Physiol.* 127 (4) (2001) 1466–1475.
- [43] M. Welch, A. Villalobos, C. Gustafsson, J. Minshull, You're one in a googol: optimizing genes for protein expression, *J. R. Soc. Interface* 6 (Suppl. 4) (2009) 467–476.
- [44] M. Welch, S. Govindarajan, J.E. Ness, A. Villalobos, A. Gurney, J. Minshull, parameters to control synthetic gene expression in *Escherichia coli*, *PLoS One* 4 (9) (2009) e7002.