

This article was downloaded by: [University of Western Cape]

On: 17 January 2012, At: 23:27

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## African Journal of Marine Science

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/tams20>

### Observations on the Diel Vertical Distribution of Hydromedusae in the Southern Benguela

E. Buecher & M. J. Gibbons

Available online: 08 Jan 2010

To cite this article: E. Buecher & M. J. Gibbons (2003): Observations on the Diel Vertical Distribution of Hydromedusae in the Southern Benguela, *African Journal of Marine Science*, 25:1, 231-238

To link to this article: <http://dx.doi.org/10.2989/18142320309504012>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.tandfonline.com/page/terms-and-conditions>

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

## OBSERVATIONS ON THE DIEL VERTICAL DISTRIBUTION OF HYDROMEDUSAE IN THE SOUTHERN BENGUELA

E. BUECHER\* and M. J. GIBBONS†

Diel vertical migration (DVM) of six common species of hydromedusae was investigated during two drogue studies conducted in St Helena Bay on the west coast of South Africa in February 1995. *Clytia* spp., *Obelia* spp. and *Bougainvillia macloviana*, were largely confined to surface waters and did not appear to display any DVM. By contrast, *Leuckartiara octona* and *Euphysa aurata* displayed clear DVM and, like *Proboscoidactyla menoni*, were found at greater depths than the other species. The depth distribution and amplitude of migration varied between surveys for most species, so definitive interpretations of patterns could not be made.

Key words: Benguela ecosystem, diel vertical migration, hydromedusa

Light, heat, nutrients and oxygen concentrations vary with depth in the oceans, as too do water motion and pressure. These changes result in a layered structure to the ocean, which organisms respond to in either a fixed (McGowan and Walker 1979) or variable manner (e.g. Ambler and Miller 1987). Although variations in vertical distribution may follow a spatial pattern (e.g. latitudinally – Vinogradov 1970), they have been better studied when they follow a temporal pattern (e.g. seasonally – Williams and Conway 1981, Buecher and Gibbons 1999; or daily – Roe *et al.* 1984). If there is a degree of spatial return to the pattern, then the variation in the depth of occupancy over a daily cycle is known as diel vertical migration (DVM). The main cue for the behaviour is thought to be light (Verwey 1966), but the causal mechanism is considered to be, among others, predatory avoidance (Gabriel and Thomas 1988). DVM has evolved independently among pelagic taxa, and is seen in all major groups, from photosynthetic micro-algae (Pitcher *et al.* 1998) to fish (Huse *et al.* 1998). The minutiae of DVM (precise depths of occupation, duration of occupancy, etc.) appear to be modified by genetic factors (Gabriel and Thomas 1988), as well as the environment (e.g. Verheye and Field 1992, Gibbons 1993).

Knowledge of the vertical distribution and DVM of plankton off southern Africa is limited, although some data are available for algae (Brown and Field 1985), chaetognaths (Gibbons 1992, Gibbons and Stuart 1994), tunicates (Gibbons 1997), copepods (Peterson *et al.* 1990) and euphausiids (Barange 1990). However, data on the vertical distribution and DVM of hydromedusae are scarce (but see the study by Pagès and Gilli

1992 in the northern Benguela). There is some evidence to suggest that DVM is displayed by the scyphomedusa *Chrysaora hysoscella* in the northern Benguela (Sparks *et al.* 2001).

Medusae are voracious predators of zooplankton (Matsakis and Conover 1991, Purcell 1992), and there is evidence to implicate them in changes of entire pelagic assemblages (Möller 1984, Purcell *et al.* 1994). Ignorance of the vertical distribution of hydromedusae impacts on the understanding of spatial dynamics of local zooplankton population. This study attempts to redress this lack of knowledge in the southern Benguela.

### MATERIAL AND METHODS

Data were collected during two drogue studies conducted in the St Helena Bay region on the west coast of South Africa. The first survey (Survey 1: 17–19 February 1995) was conducted off Cape Columbine (30°10'S, 17°44'E) in water 168 m deep, the second (Survey 2: 20–26 February 1995) in St Helena Bay (32°35'S, 18°07'E) in water 72 m deep.

Zooplankton samples were collected at regular times throughout the day and the night (02:00, 04:00, 10:30 or 12:30, and 20:30) using a 200- $\mu$ m meshed 1 m<sup>2</sup> multiple opening-closing rectangular midwater trawl (RMT-1 $\times$ 6). The trawl was towed obliquely for a short duration (<1 minute) over five depth strata (Survey 1: ~0–20, 20–40, 40–60, 60–100 and 100–150 m; Survey 2: ~0–10, 10–20, 20–30,

\* Zoology Department, University of the Western Cape, Private Bag X17, Bellville 7535, South Africa; and Marine & Coastal Management, Private Bag X2, Rogge Bay 8012, Cape Town, South Africa. E-mail: ebuecher@mcm.wcape.gov.za

† Zoology Department, University of the Western Cape




Fig. 1: Temporal variations in the vertical distribution of temperature, oxygen and Chlorophyll *a* concentration during (a) Survey 1 and (b) Survey 2.  $D_1$  denotes first day of sampling (Survey 1; 17 February 1995, Survey 2; 20 February 1995)

Table 1: Comparison of day and night total average abundances and weighed mean depth (*WMD*) for the six hydromedusae species collected in Surveys 1 and 2. Average *WMD* was calculated using only the values (depths) where the species were collected. Subscript indices in parenthesis indicate the number of times the species was collected

Species	Survey 1				Survey 2			
	Abundance (Individuals 100 m <sup>-3</sup> ± SE)		WMD (m)		Abundances (Individuals 100 m <sup>-3</sup> ± SE)		WMD (m)	
	Day n = 3	Night n = 8	Day n = 3	Night n = 8	Day n = 6	Night n = 17	Day n = 6	Night n = 17
<i>E. aurata</i>	0.4 ± 0.1 <sub>(3)</sub>	0.8 ± 0.1 <sub>(8)</sub>	125.3 ± 1.6	74.6 ± 5.4	0.03 ± 0.03 <sub>(1)</sub>	0.9 ± 0.03 <sub>(11)</sub>	2.5	33.5 ± 4.9
<i>L. octona</i> +	0.3 ± 0.1 <sub>(3)</sub>	0.1 ± 0.03 <sub>(8)</sub>	82.7 ± 16.7	61.0 ± 3.6	0.1 ± 0.07 <sub>(5)</sub> *	0.6 ± 0.1 <sub>(17)</sub>	15.9 ± 1.9	18.5 ± 2.1
<i>Chytia</i> spp.	nc	0.003 ± 0.002 <sub>(2)</sub>	nc	40.5 ± 4.8	12.3 ± 5.0 <sub>(6)</sub> >*	3.6 ± 1.2 <sub>(5)</sub>	10.7 ± 1.4	17.8 ± 2.4
<i>P. menoni</i>	0.01 <sub>(1)</sub>	0.003 ± 0.002 <sub>(2)</sub>	129.5	101.0 ± 14.7	0.5 ± 0.3 <sub>(5)</sub>	0.1 ± 0.03 <sub>(10)</sub>	20.8 ± 4.5	24.0 ± 4.1
<i>Obelia</i> spp.	nc	nc	nc	nc	2.0 ± 0.7 <sub>(5)</sub>	1.4 ± 0.6 <sub>(9)</sub>	15.3 ± 2.0	15.0 ± 1.0
<i>B. macloviana</i> +	nc	nc	nc	nc	4.3 ± 1.5 <sub>(6)</sub>	3.4 ± 1.0 <sub>(17)</sub>	5.2 ± 0.3	15.2 ± 3.1

nc = Not collected

+ = Presence of ocelli

\* Significant differences ( $\alpha = 0.05$ )

30–40 and 40–60 m). Flow rates were estimated using a flowmeter mounted above the mouth of the net, and 100% efficiency was assumed for each tow. All zooplankton samples were preserved in 4% buffered saline formalin.

Temperature, salinity, dissolved oxygen and chlorophyll concentration were measured (over depth) at regular times (04:00, 09:00, 18:00 and 22:00) throughout the study. The methods used are detailed in Kiørboe *et al.* (1998) and Gibbons *et al.* (1999).

All medusae from the samples were counted, and the stratified density in each depth stratum (individuals 100 m<sup>-3</sup>) was calculated from knowledge of the volume filtered by the nets. For every each sampling time, the average abundance (calculated using the abundances in each of the different depth strata) and the weighted mean depths (*WMD*; see Roe *et al.* 1984) of the populations were calculated. The extent of DVM was estimated (when possible) as the maximum distance covered between consecutive day-night estimates of *WMD*. Day and night differences between *WMD* were compared using Student's *t*-test.

## RESULTS

During the study period, sunrise was between 06:22 and 06:31 and sunset between 19:27 and 19:38. Detailed descriptions of the hydrological conditions during the surveys are given in Kiørboe *et al.* (1998) and Gibbons *et al.* (1999) and are briefly summarized here.

There was warming of the surface waters during Survey 1 (Fig. 1), causing a deepening of the thermocline (marked by the 10°C isotherm) from 65 (Day 1) to 80 m (Day 2). The water column was isohaline (34.7) throughout. Concentrations of dissolved oxygen and chlorophyll *a* increased in the upper 20 m after Day 1, peaking at 7.21 ml l<sup>-1</sup> and 14.09 mg m<sup>-3</sup> respectively on Day 3.

In Survey 2, there was an increase in both surface temperature (12 to 14.7°C) and the thermal gradient (3.1 to 5.5°C) over the study period. The temperature was stable (9.4°C) below the thermocline, which was situated between 35 and 40 m. There were diurnal variations in surface temperature, especially at the end of the survey. The water column was isohaline throughout (34.7). The upper layers (>20 m) was well oxygenated (mean 5.8 ml l<sup>-1</sup>), but the oxygen concentration decreased markedly to <1 ml l<sup>-1</sup> below 30 m. Highest chlorophyll *a* concentrations were found in the upper 20 m.

The results given here cover six of the commonest hydromedusae collected. Two species of Anthomedusae (*Euphysa aurata* and *Leuckartiara octona*) and Lepto-

(a) Survey 1

(b) Survey 2




Fig. 2: Temporal changes in the weighted mean depth (*WMD*) and the average abundances of medusae during (a) Survey 1 and (b) Survey 2. The shaded areas are night-time periods and the dotted line indicates the position of the thermocline

medusae (*Clytia* spp. and *Proboscidactyla menoni*) were found in both surveys, and an additional two species, *Bougainvillia macloviana* (Anthomedusae) and *Obelia* spp. (Leptomedusae) were collected during Survey 2. No temporal succession of the different species was noted during either survey. Medusae were most abundant during Survey 2, except for *E. aurata* (Fig. 2).

The most abundant species during Survey 1 was *E. aurata* (Fig. 2), and twice as many individuals were collected at night than during the day (Table I). Maximum abundance averaged 1.6 individuals 100 m<sup>-3</sup>. *E. aurata* was shallower during the early part of the night than during the day. Its maximum vertical range was 71 m, corresponding to 42% of the water column. *E. aurata* was less abundant on Survey 2 and it was rarely caught during the day, which makes day-night comparisons of either abundance or distribution difficult. The species was found throughout the water column and appeared to cross the thermocline.

The abundance of *L. octona* varied throughout Survey 1, with no clear diel differences. The species was shallower at night. Because most of the population remained at 60–70 m throughout the survey, the amplitude of migration was difficult to determine. However, it appears that the species can traverse a depth range of 70 m. In contrast, during Survey 2, *L. octona* were more abundant at night than during the day, being restricted to the upper 30 m and above the thermocline.

*Clytia* spp. was recorded only twice during Survey 1, between 30 and 50 m deep. However, it was the most abundant medusa found during Survey 2, attaining a maximum average abundance of 32.19 individuals 100 m<sup>-3</sup>. Average abundances differed significantly ( $p < 0.05$ ) between night and day during Survey 2 and remained in the upper 30 m throughout.

*P. menoni* were collected three times during Survey 1, between 70 and 130 m deep. During Survey 2, they were collected throughout the water column, but tended to occupy mid-depths. There was no discernible diel differences in abundance.

The abundances of *Obelia* spp. and *B. macloviana* during the latter stages of Survey 2. Although there is some evidence to suggest that abundances were highest at night, this was not consistent. The populations of both species were centred in the upper 20 m and DVM was not observed.

## DISCUSSION

All the species under study have a global distribution, except in the polar areas (Kramp 1961), and have been

collected regularly during previous surveys in the Benguela (Pagès *et al.* 1992, Buecher and Gibbons 2000). Although the overall diversity and the abundance of the medusae were both lower than previously reported in the region (Buecher and Gibbons 2000), this reflects the small number of samples collected, as well as their restricted geographic coverage. For Survey 1, it might also reflect station location, because areas at or close to upwelling regions are believed to have a depauperate (and certainly different) zooplankton assemblage relative to areas downstream (Gibbons and Hutchings 1996). The abundance of medusae (especially meroplanktonic species, as observed here) is extremely variable (Buecher and Gibbons 2000), and reflects current and historic physical processes that result in changes to the autochthonous and allochthonous composition of assemblages.

In the absence of appreciable vertical mixing or advection during the study period (also Gibbons *et al.* 1999), changes in the abundance of medusae linked to the diel cycle (e.g. in the case of *Clytia* spp. or *Euphysa aurata*) can possibly be attributed to recruitment to the water column from deep water, given that the layers above and below the thermocline were moving in slightly different directions (see Gibbons *et al.* 1999). A diel signal to abundance complicates any clear interpretation of DVM, and consequently, discussion is confined here to only those species that showed consistent or unambiguous patterns of vertical distribution.

Three species, *Clytia* spp., *Obelia* spp. and *B. macloviana*, which were common during Survey 2, were largely confined to the upper layers of the water column and did not appear to display any DVM. Such meroplanktonic species are more common in shallow than in deep water (Moreira 1973) and typically occur above the thermocline. Although several species of *Clytia*, *Obelia* and *Bougainvillia* have on occasions failed to display DVM (Russell 1931, Palma 1985, Ballard and Myers 1996), there are instances when such vertical behaviour has been clear (Russell 1931, Mills 1981). From three different surveys of various duration (1–4 nights) conducted annually between 1924 and 1926, Russell (1931) showed that some species of medusae, especially shallow-water ones, behaved differently from one year to the next. These changes in behaviour are not thought to be intrinsic (Mills 1981) and seem to be controlled mainly by the light intensity (day or moonlight), which also influences the timing of the migrations (e.g. downward migration at night or dawn). If these migrations vary from one year to the next, they can also differ from one season to another (Moreira 1973).

DVM was not clear in *P. menoni* (during either survey), and *L. octona* and *E. aurata* migrated during

Table II: Comparison of vertical migrations from various species of the same genus elsewhere with those observed in the present study. DVM means the population performs an upward migration at night and has a deeper distribution during the day. Reverse DVM means the population has a distribution close to the surface during the day and perform a downward migration at night

Species	Behaviour	Location	Source
<i>Euphysa aurata</i>	DVM	Villefranche Bay, France	Palma (1985)
<i>Leuckartiara octona</i>	DVM	English Channel, UK	Russell (1931)
<i>Clytia simplex</i>	No migration	Benguela current, Namibia	Pagès and Gili (1992)
<i>Phialidium gregarium</i>	Reverse DVM*	Friday Harbor, USA	Mills (1981)
	Reverse DVM	Saanich Inlet, USA	
<i>Phialidium</i> spp.	No migration	Villefranche Bay, France	Palma (1985)
<i>Phialidium</i> spp.	Observation of some upward movement at dusk, but no clear DVM		
<i>Proboscidactyla ornata</i>	DVM	English Channel, UK	Russell (1931)
<i>Obelia</i> spp.	DVM	Coast of Santos, Brazil	Moreira (1973)
<i>Obelia</i> spp.	DVM	Coast of Santos, Brazil	Moreira (1973)
<i>Obelia</i> spp.	No clear DVM	English Channel, UK	Gough (1905)
<i>Obelia</i> spp.	No clear DVM	English Channel, UK	Russell (1931)
<i>Bougainvillia</i> spp.	DVM*	Friday Harbor, USA	Mills (1981)
<i>Bougainvillia</i> spp.	DVM	Plymouth, UK	Russell (1928)
<i>B. britannica</i> , <i>B. principis</i> , <i>B. ramosa</i> , <i>B. pyramidata</i>	No migration	Lough Hyne, Ireland	Ballard and Myers (1996)

\* Laboratory observations

Survey 1 but not on Survey 2. Species from these three genera have been shown to display DVM (Russell 1931, Moreira 1973, Palma 1985; Table II). However, the species in the present surveys migrated deeper during Surveys 1 and 2 than the depths attained by the species in the aforementioned studies. Russell (1931) observed that macrozooplankton in deep water display clearer patterns of DVM than those in shallow water. Also, Andersen *et al.* (1997) noted that the amplitude of migration of species occupying shallow water was smaller relative to those occupying deeper water. Although *P. menoni* and *L. octona* crossed the thermocline during Survey 2, this boundary limited the vertical migration of the former species in southern Brazil (Moreira 1973).

There was no difference in this study between Antho- or Leptomedusae regarding their DVM or vertical distribution. In contrast, Costello and Colin (1995) suggested that bullet-shaped species (principally Anthomedusae) might be more active than the dish-shaped species, because of their better jet propulsion. Also, there was no relationship between DVM and the presence of light-sensitive ocelli in the species under study (Table I). Although ocelli have been shown to play a role in DVM of some medusae species (Mills 1983), other studies have found that ocelli were not always present in light-sensitive migrating species (Moreira 1973, Mills and Goy 1988). However, some photo-sensitive neurons have been recognized as responsible for a swimming reaction when stimulated by light (Anderson and Mackie 1977), which can then initiate

migratory behaviour.

Despite the present inconclusive results (largely on account of low animal densities), to the authors' knowledge the present data represent the longest published time-series describing DVM of Hydromedusae. The discrepancies and inconsistencies in the patterns found here over three and six-day surveys emphasize the fact that 24 h of observations are not adequate to describe the migrating behaviour of medusae definitively. The fact that the present observations are contradictory and sometimes contrast with previous observations show the lack of understanding about DVM of the species under study. For example, Mills (1981) suggested that DVM of medusae allows individuals to concentrate close to the surface for reproductive purposes. However, such factors as presence of prey, pressure changes within the water column or diel differences in buoyancy need to be taken into consideration to explain vertical distribution of medusae (Arai 1992). Further observations are clearly necessary to understand this migratory behaviour and the biotic and abiotic factors that control it.

#### ACKNOWLEDGEMENTS

We thank the officers and crew of the F. R. S. *Africana* and our colleagues for their co-operation and assistance at sea. The National Research Foundation is gratefully acknowledged for financial support.

## LITERATURE CITED

- AMBLER, J. W. and C. B. MILLER 1987 — Vertical habitat-partitioning by copepodites and adults of subtropical oceanic copepods. *Mar. Biol.* **94**(4): 561–577.
- ANDERSEN, V., SARDOU, J. and B. GASSER 1997 — Macroplankton and micronekton in the northeast tropical Atlantic: abundance, community composition and vertical distribution in relation to different trophic environments. *Deep-Sea Res.* **44**: 193–222.
- ANDERSON, P. A. V. and G. O. MACKIE 1977 — Electrically coupled, photosensitive neurons control swimming in a jellyfish. *Science* **197**: 186–188.
- ARAI, M. N. 1992 — Active and passive factors affecting aggregations of Hydromedusae: a review. In *Aspects of Hydrozoan Biology*. Bouillon, J., Boero, F., Cicogna, F., Gili, J.-M. and R. G. Hughes (Eds). *Scientia Mar., Barcelona* **56**: 99–108.
- BALLARD, L. and A. MYERS 1996 — Seasonal changes in the vertical distribution of five species of the family Bougainvillidae (Cnidaria : Anthomedusae) at Lough Hyne, southwest Ireland. In *Advances in Hydrozoan Biology*. Piraino, S., Boero, F., Bouillon, J., Cornelius, P. F. S. and J.-M. Gili (Eds). *Scientia Mar., Barcelona* **60**: 69–74.
- BARANGE, M. 1990 — Vertical migration and habitat partitioning of six euphausiid species in the northern Benguela upwelling system. *J. Plankt. Res.* **12**(6): 1223–1237.
- BROWN, P. C. and J. G. FIELD 1985 — Diel variation in production rates of natural phytoplankton populations in the southern Benguela upwelling region. *Botanica mar.* **28**: 201–208.
- BUECHER, E. and M. J. GIBBONS 1999 — Temporal persistence in the vertical structure of the assemblage of planktonic medusae in the NW Mediterranean Sea. *Mar. Ecol. Prog. Ser.* **189**: 105–115.
- BUECHER, E. and M. J. GIBBONS 2000 — Interannual variation in the composition of the assemblages of Medusae and Ctenophores in St Helena Bay, southern Benguela ecosystem. In *Trends in Hydrozoan Biology*. 4. Mills, C. E., Boero, F., Migotto, A. and J.-M. Gili (Eds). *Scientia Mar., Barcelona* **64**: 123–134.
- COSTELO, J. H. and S. P. COLIN 1995 — Flow and feeding by swimming scyphomedusae. *Mar. Biol.* **124**: 399–406.
- GABRIEL, W. and B. THOMAS 1988 — Ultimate causes of vertical migration in zooplankton: an evaluation by evolutionary game theory. In *Ecodynamics. Contributions to Theoretical Ecology*. Wolff, W., Soeden, C. J. and F. R. Dreppen (Eds). Berlin; Springer: 127–134.
- GIBBONS, M. J. 1992 — Diel feeding and vertical migration of *Sagitta serratodentata* Krohn *tasmanica* Thomson (Chaetognatha) in the southern Benguela. *J. Plankt. Res.* **14**(2): 249–259.
- GIBBONS, M. J. 1993 — Vertical migration and feeding of *Euphausia lucens* at two 72 h stations in the southern Benguela upwelling region. *Mar. Biol.* **116**(2): 257–268.
- GIBBONS, M. J. 1997 — Feeding and migration of *Thalia democratica* on the Agulhas Bank, South Africa. *J. mar. biol. Ass. U.K.* **77**: 493–505.
- GIBBONS, M. J. and V. STUART 1994 — Feeding and vertical migration of the chaetognath *Sagitta friderici* (Ritter-Zahony, 1911) in the southern Benguela during spring 1987, with notes on seasonal variability of feeding ecology. *S. Afr. J. mar. Sci.* **14**: 361–372.
- GIBBONS, M. J. and L. HUTCHINGS 1996 — Zooplankton diversity and community structure around southern Africa, with special attention to the Benguela upwelling system. *S. Afr. J. Sci.* **92**: 63–76.
- GIBBONS, M. J., GUGUSHE, N., BOYD, A. J., SHANNON, L. J. and B. A. MITCHELL-INNES 1999 — Changes in the composition of the non-copepod zooplankton assemblage in St Helena Bay (southern Benguela ecosystem) during a six-day drogue study. *Mar. Ecol. Prog. Ser.* **180**: 111–120.
- GOUGH, L. H. 1905 — On the distribution and migrations of *Muggiaea atlantica* Cunningham, in the English Channel, the Irish Sea and off the South and West Coasts of Ireland in 1904. *J. Cons. perm. int. Explor. Mer. Publ. Cir.* **29**: 493–519.
- HUSE, I., HAMUKUAYA, H., BOYER, D. C., MALAN, P. E. and T. STRØMME 1998 — The diurnal vertical dynamics of Cape hakes and their potential prey. In *Benguela Dynamics: Impacts of Variability on Shelf-Sea Environments and their Living Resources*. Pillar, S. C., Moloney, C. L., Payne, A. I. L. and F. A. Shillington (Eds). *S. Afr. J. mar. Sci.* **19**: 365–376.
- KIØRBOE, T., TISELIUS, P., MITCHELL-INNES, B. A., HANSEN, J. L. S., VISSER, A. W. and X. MARI 1998 — Intensive aggregate formation with low vertical flux during an upwelling-induced diatom bloom. *Limnol. Oceanogr.* **43**: 104–116.
- KRAMP, P. L. 1961 — Synopsis of the medusae of the world. *J. mar. biol. Ass. U.K.* **40**: 1–469.
- MATSAKIS, S. and R. J. CONOVER 1991 — Abundance and feeding of medusae and their potential impact as predators on other zooplankton in Bedford Basin (Nova Scotia, Canada) during spring. *Can. J. Fish. aquat. Sci.* **48**: 1419–1430.
- MCGOWAN, J. A. and P. W. WALKER 1979 — Structure in the copepod community of the north Pacific central gyre. *Ecol. Monogr.* **49**(2): 195–226.
- MILLS, C. E. 1981 — Diversity of swimming behaviors in hydromedusae as related to feeding and utilization of space. *Mar. Biol.* **64**(2): 184–189.
- MILLS, C. E. 1983 — Vertical migration and diel activity patterns of Hydromedusae: studies in a large tank. *J. Plankt. Res.* **5**: 619–635.
- MILLS, C. E. and J. GOY 1988 — *In situ* observations of the behavior of mesopelagic *Solmissus* Narcomedusae (Cnidaria, Hydrozoa). *Bull. mar. Sci.* **43**: 739–751.
- MOREIRA, G. S. 1973 — On the diurnal vertical migration of Hydromedusae off Santos, Brazil. In *Recent Trends in Research in Cœlenterate Biology*. Tokiota, T. and T. Nishimura (Eds). *Publs Seto mar. biol. Lab.* **20**: 537–566.
- MÖLLER, H. 1984 — Reduction of a larval herring population by jellyfish predator. *Science, N.Y.* **224**(4649): 621–622.
- PAGÈS, F., GILI, J.-M. and J. BOUILLON 1992 — Medusae (Hydrozoa, Scyphozoa, Cubozoa) of the Benguela Current (South eastern Atlantic). *Scientia Mar., Barcelona* **56**: 1–64.
- PAGÈS, F. and J.-M. GILI 1992 — Influence of the thermocline on the diel migration of medusae during a 48 h sampling period. *S. Afr. J. Zool.* **27**: 50–59.
- PALMA, S. G. 1985 — Migración nictemeral del macroplankton gelatinoso de la bahía de Villefranche-sur-Mer, Mediterráneo Noroccidental. *Investigación pesq., Barcelona* **49**: 261–274.
- PETERSON, W. T., PAINTING, S. J. and L. HUTCHINGS 1990 — Diel variations in gut pigment content, diel vertical migration and estimates of grazing impact for copepods in the southern Benguela upwelling region in October 1987. *J. Plankt. Res.* **12**(2): 259–281.
- PITCHER, G. C., BOYD, A. J., HORSTMAN, D. A. and B. A. MITCHELL-INNES 1998 — Subsurface dinoflagellate populations, frontal blooms and the formation of red tide in the southern Benguela upwelling system. *Mar. Ecol. Prog. Ser.* **172**: 253–264.
- PURCELL, J. E. 1992 — Effects of predation by the scyphomedusan *Chrysaora quinquecirrha* on zooplankton populations in Chesapeake Bay, USA. *Mar. Ecol. Prog. Ser.* **87**: 65–76.
- PURCELL, J. E., WHITE, J. R. and M. R. ROMAN 1994 —



- Predation by gelatinous zooplankton and resource limitation as potential controls of *Acartia tonsa* copepod populations in Chesapeake Bay. *Limnol. Oceanogr.* **39**: 263–278.
- ROE, H. S. J., ANGEL, M. V., BADCOCK, J., DOMANSKI, P., JAMES, P. T., PUGH, P. R. and M. H. THURSTON 1984 — The diel migrations and distributions within a mesopelagic community in the North East Atlantic. 1. Introduction and sampling procedures. *Prog. Oceanogr.* **13**: 245–268.
- RUSSELL, F. S. 1928 — The vertical distribution of marine macroplankton. 4. Further observation on diurnal changes. *J. mar. biol. Ass. U.K.* **15**: 81–103.
- RUSSELL, F. S. 1931 — The vertical distribution of marine macroplankton. 11. Further observations on diurnal changes. *J. mar. biol. Ass. U.K.* **17**: 767–784.
- SPARKS, C., BUECHER, E., BRIELEY, A. S., AXELSEN, B. E., BOYER, H. and M. J. GIBBONS 2001 — Observations on the distribution, and relative-abundance of *Chrysaora hysoscella* (Cnidaria, Scyphozoa) and *Aequorea aequorea* (Cnidaria, Hydrozoa) in the northern Benguela ecosystem. *Hydrobiologia* **451**: 275–286.
- VERHEYE, H. M. and J. G. FIELD 1992 — Vertical distribution and diel vertical migration of *Calanoides carinatus* (Krøyer, 1849) developmental stages in the southern Benguela upwelling region. *J. expl mar. Biol. Ecol.* **158**: 123–140.
- VERWEY, J. 1966 — The role of some external factors in the vertical migration of marine animals. *Neth. J. Sea Res.* **3**: 245–266.
- VINOGRADOV, M. E. 1970 — *Vertical Distribution of the Oceanic Zooplankton*. Jerusalem; Israel Program for Scientific Translations: 339 pp.
- WILLIAMS, R. B. and D. V. P. CONWAY 1981 — Vertical distribution and seasonal abundance of *Aglantha digitale* (O. F. Müller) (Coelentera: Trachymedusae) and other planktonic coelenterates in the Northeast Atlantic Ocean. *J. Plankt. Res.* **3**: 633–643.