VIROLOGY DIVISION NEWS



Bacterial Viruses Subcommittee and Archaeal Viruses Subcommittee of the ICTV: update of taxonomy changes in 2021

Mart Krupovic¹® · Dann Turner²® · Vera Morozova³® · Mike Dyall-Smith⁴® · Hanna M. Oksanen⁵® · Rob Edwards⁶® · Bas E. Dutilh¹® · Susan M. Lehman⁶® · Alejandro Reyesഐ · Diana P. Baquero¹® · Matthew B. Sullivan¹⁰,¹¹® · Jumpei Uchiyama¹²® · Jesca Nakavuma¹³® · Jakub Barylski¹⁴ · Mark J. Young¹⁵® · Shishen Du¹⁶® · Poliane Alfenas-Zerbini¹¹® · Alla Kushkina¹⁶® · Andrew M. Kropinski¹⁰,²⁰® · Ipek Kurtböke²¹® · J. Rodney Brister²² · Cédric Lood²³® · B. L. Sarkar²⁴ · Tong Yigang²⁵® · Ying Liu¹ · Li Huang²⁶® · Johannes Wittmann²γ · Nina Chanishvili²⁶ · Leonardo J. van Zyl²⁰® · Janis Rumnieks³⁰® · Tomohiro Mochizuki³¹® · Matti Jalasvuori³²® · Ramy K. Aziz³³® · Małgorzata Łobocka³⁴® · Kenneth M. Stedman³⁵® · Andrey N. Shkoporov³⁶® · Annika Gillis³७® · Xu Peng³⁶® · François Enault³ഐ · Petar Knezevic⁴⁰® · Rob Lavigne²³® · Sung-Keun Rhee⁴¹® · Virginija Cvirkaite-Krupovic¹® · Cristina Moraru⁴²® · Andrea I. Moreno Switt⁴³® · Minna M. Poranen⁵ · Andrew Millard⁴⁴® · David Prangishvili⁴⁵ · Evelien M. Adriaenssens⁴®

© The Author(s) 2021

Abstract

In this article, we – the Bacterial Viruses Subcommittee and the Archaeal Viruses Subcommittee of the International Committee on Taxonomy of Viruses (ICTV) – summarise the results of our activities for the period March 2020 – March 2021. We report the division of the former Bacterial and Archaeal Viruses Subcommittee in two separate Subcommittees, welcome new members, a new Subcommittee Chair and Vice Chair, and give an overview of the new taxa that were proposed in 2020, approved by the Executive Committee and ratified by vote in 2021. In particular, a new realm, three orders, 15 families, 31 subfamilies, 734 genera and 1845 species were newly created or redefined (moved/promoted).

Changes in Subcommittee structure and membership

Bacteriophage and archaeal virus taxonomy has been formally under the auspices of the ICTV Bacterial and Archaeal Viruses Subcommittee, which, at its inception in 1966, was named the Viruses of Prokaryotes Subcommittee, led by David E. Bradley (https://talk.ictvonline.org/information/w/ictv-history). Given the revived interest in bacterial and archaeal viruses and recent enormous increase in the number of characterized isolates and need for creation of numerous taxa to classify them, the Executive Committee (EC) voted on the creation of two separate Subcommittees (EC51, July 2019), formally starting their mandates after EC52 (October

Handling Editor: Sead Sabanadzovic.

Published online: 21 August 2021

Evelien M. Adriaenssens evelien.adriaenssens@quadram.ac.uk

Extended author information available on the last page of the article

2020). The new Bacterial Viruses Subcommittee is chaired by Evelien Adriaenssens, supported by Dann Turner as the Vice Chair, and the new Archaeal Viruses Subcommittee is chaired by Mart Krupovic. Both Chairs were elected for three-year terms ending in 2023. As a result, this taxonomy update summarises both bacterial and archaeal virus proposals for the last time, reflecting the fact that proposals were submitted prior to the formal reorganisation of the original Subcommittee.

In the new Bacterial Viruses Subcommittee, we continue our structure of Study Groups (SGs), regional representatives and general members. We would like to welcome new members Jesca Nakavuma (Uganda), Alejandro Reyes (Colombia), Cristina Moraru (Germany), Susan Lehman (USA), Cédric Lood (Belgium) and Andrey Shkoporov (Ireland) and would like to thank those who have left the Subcommittee for their service.

In the framework of the Bacterial and Archaeal Viruses Subcommittee, all taxonomic proposals dealing with archaeal viruses were handled by a single SG. The new



Archaeal Viruses Subcommittee currently includes 11 SGs (Table 1), each created for a three-year term. Of note, the *Halopanivirales* SG, chaired by Hanna M. Oksanen (Finland), oversees the taxonomy of evolutionarily related viruses of the order *Halopanivirales*, which includes two families of viruses infecting archaea (*Sphaerolipoviridae* and *Simuloviridae*) and one family of bacteriophages (*Matsushitaviridae*). Thus, the latter SG bridges the two Subcommittees. Additional SGs will be created in the near future, in particular, to develop the taxonomy of archaeal members of the class *Caudoviricetes*; only a handful of isolated

representatives of this highly diverse and expansive group of archaeal viruses are currently classified.

Taxonomy update

We had a record year of submissions of taxonomy proposals in 2020, with 188 proposals submitted, of which all but one were approved and ratified (Supplementary Table S1). The changes at each of the taxonomic ranks in use for bacterial and archaeal viruses are summarised in Table 2. We

Table 1 Composition of the archaeal viruses subcommittee

Study Group	Member	Country		
Bicaudaviridae SG	Mart Krupovic*	France		
	Li Huang	China		
	Mark J. Young	USA		
	Virginija Cvirkaite-Krupovic	France		
Desulfurococcales viruses SG	Tomohiro Mochizuki*	Japan		
	Mart Krupovic	France		
Fuselloviridae SG	Kenneth M. Stedman*	USA		
	Mart Krupovic	France		
Halopanivirales SG	Hanna M. Oksanen*	Finland		
	Mike Dyall-Smith	Australia		
	Shishen Du	China		
	Matti Jalasvuori	Finland		
Halspiviridae SG	Mike Dyall-Smith*	Australia		
	Hanna M. Oksanen	Finland		
Ovaliviridae SG	Li Huang*	China		
Pleolipoviridae SG	Hanna M. Oksanen*	Finland		
	Mike Dyall-Smith	Australia		
	Ying Liu	France		
Portogloboviridae SG	Mart Krupovic*	France		
	Ying Liu	France		
Thaspiviridae SG	Sung-Keun Rhee*	Republic of Korea		
	Mart Krupovic	France		
Tokiviricetes SG	Mart Krupovic*	France		
	Tomohiro Mochizuki	Japan		
	Xu Peng	Denmark		
	Diana P. Baquero	France		
Turriviridae SG	Mark J. Young*	USA		

^{*}Study Group Chair

Table 2 Summary of taxonomic changes for bacterial and archaeal viruses for Master Species List 36, ratified March 2021

	Species	Genus	Subfamily	Family	Order	Class	Phylum	Kingdom	Realm
Abolished	20	2	0	0	0	0	0	0	0
New	1775	700	28	14	2	1	1	1	1
Moved or pro- moted (and/or renamed)	70	34	3	1	1	0	0	0	0
Renamed	33	8	0	1	1	1	0	0	0



created a record 1775 new species, 700 new genera, 28 new subfamilies, 14 new families and two new orders and also created one new realm containing one new kingdom, one new phylum and one new class. Given the large numbers of proposals and taxa that were created, moved or renamed, it becomes unfeasible to describe all the changes in detail; however, we urge interested readers to consult Supplementary Table S1 and the associated proposals from the ICTV website (https://talk.ictvonline.org/files/ictv_official_taxon omy_updates_since_the_8th_report/m/prokaryote-official). Instead, below we provide a brief overview of the most notable changes.

The new realm *Adnaviria*, a new megataxonomy of archaeal filamentous viruses

Recently, the taxonomic framework of the ICTV has been expanded from five to 15 ranks, with the highest-level rank, realm, being equivalent to the domain rank used for cellular organisms [1]. Until recently, four such realms had been established for classification of viruses infecting hosts from different domains of life [2, 3]. This year, a new realm, Adnaviria, was created for classification of archaeal filamentous viruses with dsDNA genomes that adopt the A-form conformation within their virions [4, 5]. The realm includes three families, Tristromaviridae, Rudiviridae and Lipothrixviridae, which are evolutionarily related to each other, but not to any other known group of viruses. The families Rudiviridae and Lipothrixviridae were already grouped together in the order Ligamenvirales [6], and a single-family order, Primavirales, has now been created for the family Tristromaviridae. The two orders are unified in the class Tokiviricetes, kingdom Zilligvirae and realm Adnaviria.

Class Caudoviricetes, order Caudovirales

We delineated three new families of short-tailed phages with small genomes that were formerly assigned to the family *Podoviridae*. The family *Salasmaviridae*, named in honour of Margarita Salas Falgueras, comprises the existing subfamily *Picovirinae*, which includes the classical bacillus phage φ29 and a range of related *Bacillus*-infecting phages with genomes between 18 and 27 kb in size. The family *Rountreeviridae*, named in honour of Phyllis Margaret Rountree, groups *Enterococcus*-infecting phages with genomes between 17 and 19 kb in size, whereas the family *Guelinviridae*, named after Antonina Guelin, groups *Clostridium*-infecting phages with genome sizes between 16 and 19 kb.

The new family *Schitoviridae*, named after Giancarlo Schito, is the formalisation of the group of N4-like phages

defined by the presence of a large virion-associated RNA polymerase, described in more detail in reference [7].

The new family *Zobellviridae*, named after Claude Zobell, groups a set of globally distributed podoviruses associated with marine ecosystems first proposed by Bischoff and colleagues [8].

Class Leviviricetes

Based on the investigation by Callanan and colleagues on the expansion of known ssRNA virus genomes [9], the family *Leviviridae* was elevated to the rank of class, named *Leviviricetes* (replacing the class *Allassoviricetes*), and expanded to include two orders (*Norzivirales* and *Timlovirales*) and six new families: *Fiersviridae* (renamed from the original family *Leviviridae*), *Atkinsviridae*, *Duinviridae*, *Solspiviridae*, *Blumeviridae* and *Steitzviridae*. A detailed description of the new taxa will be published separately.

Class Tectiliviricetes, new family Autolykiviridae

The new family *Autolykiviridae* formalises the group of nontailed dsDNA bacteriophages discovered and described by Kauffmann and colleagues [10], combining features of both corticoviruses and tectiviruses. This new family contains two new genera and five new species.

Order Halopanivirales, new families Matsushitaviridae and Simuloviridae

Until recently, the family Sphaerolipoviridae, which includes icosahedral tailless viruses with internal membranes, consisted of three genera, Alphasphaerolipovirus, Betasphaerolipovirus and Gammasphaerolipovirus. The first two of these genera included viruses infecting halophilic archaea, whereas the last one included phages infecting thermophilic bacteria [11]. Although viruses from the three genera are evolutionarily related [12], they display considerable sequence divergence. Thus, the genera Betasphaerolipovirus and Gammasphaerolipovirus have been renamed and moved from the Sphaerolipoviridae into new families, Simuloviridae and Matsushitaviridae (named after Isao Matsushita), respectively. The order Halopanivirales now contains a family of bacterial viruses, Matsushitaviridae, and two families of archaeal viruses, family Sphaerolipoviridae and Simuloviridae. As mentioned above, the order is under purview of a single Study Group, which is part of both the Archaeal Viruses SC and the Bacterial Viruses SC.



Order *Tubulavirales*, new family *Paulinoviridae*

The new family *Paulinoviridae* addresses the challenge of defining family demarcation criteria for phages with small genomes, where members of the same family would share at least two orthologous proteins. Informed by prior work on a gene-content network of predicted filamentous prophage sequences [13], we moved the genera *Bifilivirus* and *Thomixvirus* from the family *Inoviridae* to the family *Paulinoviridae*.

Online (10th) Report of the ICTV

Virus Taxonomy: The Classification and Nomenclature of Viruses - The Online (10th) Report of the ICTV is freely accessible at http://ictv.global/report, and summaries of the chapters on each virus family are published in the Journal of General Virology. In 2020, four new chapters on bacterial and archaeal viruses were produced by members of the Archaeal Viruses SC and Bacterial Viruses SC, namely, on the families *Herelleviridae* [14], *Spiraviridae* [15], *Ovaliviridae* [16] and *Finnlakeviridae* [17].

Conclusion


This past year has been extremely productive in terms of new bacterial and archaeal virus taxa described. It would not have been possible without an active pool of scientists, both within and outside the subcommittee and its study groups, who scour databases, perform analyses and submit proposals. We continue to encourage people to contact us to formalise new discoveries into the taxonomic framework and will keep reaching out to interested parties. Finally, we would like to acknowledge one person in particular, Prof Andrew Kropinski, the Subcommittee Chair from 2014-2020, who authored, co-authored and/or assisted with the majority of proposals that have been approved in the last decade.

Supplementary Information The online version contains supplementary material available at https://doi.org/10.1007/s00705-021-05205-9.

Acknowledgements We would like to acknowledge the invaluable help from colleagues at NCBI, in particular Igor Tolstoy.

Author contributions All authors contributed to the creation or assessment of taxonomy proposals; M.K. and E.M.A wrote the manuscript with input from D.T. and A.M.K; all authors evaluated and approved the final version.

Funding V.M. was supported by Russian Ministry of Education and Science Project No. 0245-2021-0008; H.M.O. was supported



Declarations

Conflict of interest All authors are current or former members of the Bacterial and Archaeal Viruses Subcommittee, Bacterial Viruses Subcommittee and/or Archaeal Viruses Subcommittee of the International Committee on Taxonomy of Viruses (ICTV).

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.



References

- Gorbalenya AE, Krupovic M, Mushegian A et al (2020) The new scope of virus taxonomy: partitioning the virosphere into 15 hierarchical ranks. Nat Microbiol 5:668–674. https://doi.org/10.1038/ s41564-020-0709-x
- Koonin EV, Dolja VV, Krupovic M et al (2020) Global organization and proposed megataxonomy of the virus world. Microbiol Mol Biol Rev 84:1–33. https://doi.org/10.1128/MMBR.00061-19
- Walker PJ, Siddell SG, Lefkowitz EJ et al (2020) Changes to virus taxonomy and the Statutes ratified by the International Committee on Taxonomy of Viruses (2020). Arch Virol 165:2737–2748. https://doi.org/10.1007/s00705-020-04752-x
- Baquero DP, Liu Y, Wang F et al (2020) Structure and assembly of archaeal viruses. Adv Virus Res 108:127–164
- Krupovic M, Kuhn JH, Wang F et al (2021) Adnaviria: a new realm for archaeal filamentous viruses with linear A-form double-stranded DNA genomes. J Virol. https://doi.org/10.1128/JVI. 00673-21
- Prangishvili D, Krupovic M (2012) A new proposed taxon for double-stranded DNA viruses, the order "Ligamenvirales." Arch Virol 157:791–795. https://doi.org/10.1007/s00705-012-1229-7
- Wittmann J, Turner D, Millard A et al (2020) From orphan phage to a proposed new family—the diversity of N4-like viruses. Antibiotics 9:663. https://doi.org/10.3390/antibiotics9100663
- Bischoff V, Bunk B, Meier-Kolthoff JP et al (2019) Cobaviruses a new globally distributed phage group infecting Rhodobacteraceae in marine ecosystems. ISME J 13:1404–1421. https://doi. org/10.1038/s41396-019-0362-7
- Callanan J, Stockdale SR, Shkoporov A et al (2020) Expansion of known ssRNA phage genomes: from tens to over a thousand. Sci Adv 6:eaay5981. https://doi.org/10.1126/sciadv.aay5981
- Kauffman KM, Hussain FA, Yang J et al (2018) A major lineage of non-tailed dsDNA viruses as unrecognized killers of marine

- bacteria. Nature 554:118–122. https://doi.org/10.1038/nature25474
- Pawlowski A, Rissanen I, Bamford JKH et al (2014) Gammasphaerolipovirus, a newly proposed bacteriophage genus, unifies viruses of halophilic archaea and thermophilic bacteria within the novel family Sphaerolipoviridae. Arch Virol 159:1541–1554. https://doi.org/10.1007/s00705-013-1970-6
- Demina T, Pietilä M, Svirskaitė J et al (2017) HCIV-1 and other tailless icosahedral internal membrane-containing viruses of the family sphaerolipoviridae. Viruses 9:32. https://doi.org/10.3390/ v9020032
- 13. Roux S, Krupovic M, Daly RA et al (2019) Cryptic inoviruses revealed as pervasive in bacteria and archaea across Earth's biomes. Nat Microbiol 4:1895–1906. https://doi.org/10.1038/s41564-019-0510-x
- Barylski J, Kropinski AM, Alikhan N-F, Adriaenssens EM (2020) ICTV virus taxonomy profile: herelleviridae. J Gen Virol 101:3–4. https://doi.org/10.1099/jgv.0.001392
- Prangishvili D, Mochizuki T, Krupovic M (2020) ICTV virus taxonomy profile: spiraviridae. J Gen Virol 101:240–241. https:// doi.org/10.1099/jgv.0.001385
- Huang L, Wang H (2021) ICTV virus taxonomy profile: ovaliviridae. J Gen Virol. https://doi.org/10.1099/jgv.0.001546
- Mäntynen S, Laanto E, Sundberg LR et al (2020) ICTV virus taxonomy profile: Finnlakeviridae. J Gen Virol 101:894–895. https:// doi.org/10.1099/jgv.0.001488

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Authors and Affiliations

- ¹ Archaeal Virology Unit, Institut Pasteur, Paris, France
- Department of Applied Sciences, Faculty of Health and Applied Sciences, University of the West of England, Bristol, UK
- Laboratory of Molecular Microbiology, Institute of Chemical Biology and Fundamental Medicine SB RAS, Novosibirsk, Russia
- Veterinary Biosciences, Faculty of Veterinary and Agricultural Sciences, University of Melbourne, Parkville, Australia

- Molecular and Integrative Biosciences Research Programme, Faculty of Biological and Environmental Sciences, University of Helsinki, Helsinki, Finland
- Flinders Accelerator for Microbiome Exploration, Adelaide, Australia
- Theoretical Biology and Bioinformatics, Department of Biology, Utrecht University, Utrecht, Netherlands
- Center for Biologics Evaluation and Research, US Food and Drug Administration, 10903 New Hampshire Ave, Silver Spring, MD 20903, USA



- Max Planck Tandem Group in Computational Biology, Department of Biological Sciences, Universidad de Los Andes, Bogotá, Colombia
- Department of Microbiology, The Ohio State University, Columbus, OH 43210, USA
- Department Civil, Environmental, and Geodetic Engineering, The Ohio State University, Columbus, OH 43210, USA
- School of Veterinary Medicine, Azabu University, Sagamihara, Japan
- ¹³ College of Veterinary Medicine, Animal Resources and Biosecurity, Makerere University, Kampala, Uganda
- Department of Molecular Virology, Faculty of Biology, Institute of Experimental Biology, Adam Mickiewicz University in Poznan, Collegium Biologicum-Umultowska 89, 61-614 Poznan, Poland
- Department of Plant Sciences and Plant Pathology, Montana State University, Bozeman, MT, USA
- Department of Microbiology, College of Life Sciences, Wuhan University, Wuhan, China
- Department of Microbiology, Universidade Federal de Viçosa, Viçosa, Brazil
- ¹⁸ Zabolotny Institute of Microbiology and Virology, NAS of Ukraine, Kyiv, Ukraine
- Department of Food Science, University of Guelph, Guelph, Canada
- Department of Pathobiology, University of Guelph, Guelph, Canada
- ²¹ University of the Sunshine Coast, Sippy Downs, Australia
- National Center for Biotechnology Information, National Library of Medicine, National Institutes of Health, Bethesda, USA
- ²³ KU Leuven, Leuven, Belgium
- Emeritus ICMR-National Institute of Cholera and Enteric Diseases (NICED), Kolkata, India
- 25 Beijing University of Chemical Technology, Beijing, China
- State Key Laboratory of Microbial Resources, Institute of Microbiology, Chinese Academy of Sciences, Beijing, China
- ²⁷ Leibniz Institute DSMZ-German Collection of Microorganisms and Cell Cultures GmbH, Berlin, Germany

- The Eliava Institute of Bacteriophage, MIcrobiology and Virology, Tbilisi, Georgia
- University of the Western Cape, Western Cape, South Africa
- ³⁰ Latvian Biomedical Research and Study Center, Riga, Latvia
- 31 Earth-Life Science Institute, Tokyo Institute of Technology, Tokyo, Japan
- ³² University of Jyvaskyla, Jyvaskyla, Finland
- Department of Microbiology and Immunology, Faculty of Pharmacy, Cairo University and Microbiology and Immunology Research Program, Children's Cancer Hospital Egypt, Cairo 57357, Egypt
- 34 Laboratory of Bacteriophage Biology, Institute of Biochemistry and Biophysics of the Polish Academy of Sciences, Warsaw, Poland
- 35 Biology Department and Center for Life in Extreme Environments, Portland State University, Portland, USA
- ³⁶ University College Cork, Cork, Ireland
- ³⁷ Laboratory of Food and Environmental Microbiology, Earth and Life Institute, UC Louvain, Louvain-la-Neuve, Belgium
- Microbial Immunity Group, Department of Biology, University of Copenhagen, Copenhagen, Denmark
- ³⁹ Université Clermont Auvergne, CNRS, LMGE, 63000 Clermont-Ferrand, France
- Faculty of Sciences Department of Biology and Ecology, University of Novi Sad, Novi Sad, Serbia
- Department of Biological Sciences and Biotechnology, Chungbuk National University, Cheongju, South Korea
- ⁴² Institute for Chemistry and Biology of the Marine Environment, University of Oldenburg, Oldenburg, Germany
- Escuela de Medicina Veterinaria, Pontificia Universidad Católica de Chile, Santiago, Chile
- ⁴⁴ Department of Genetics and Genome Biology, University of Leicester, Leicester, UK
- ⁴⁵ Ivane Javakhishvili Tbilisi State University, Tbilisi, Georgia
- 46 Quadram Institute Biosciences, Norwich Research Park, Norwich NR4 7UQ, UK

