





REVIEW

Open Access



The impact of mitochondrial impairments on sperm function and male fertility: a systematic review

Minoo Vahedi Raad^{1†}, Amir Masoud Firouzabadi^{2,3†}, Maryam Tofighi Niaki⁴, Ralf Henkel^{5,6,7*} and Farzaneh Fesahat^{2*}

Abstract

Background Besides adenine triphosphate (ATP) production for sustaining motility, the mitochondria of sperm also host other critical cellular functions during germ cell development and fertilization including calcium homeostasis, generation of reactive oxygen species (ROS), apoptosis, and in some cases steroid hormone biosynthesis. Normal mitochondrial membrane potential with optimal mitochondrial performance is essential for sperm motility, capacitation, acrosome reaction, and DNA integrity.

Results Defects in the sperm mitochondrial function can severely harm the fertility potential of males. The role of sperm mitochondria in fertilization and its final fate after fertilization is still controversial. Here, we review the current knowledge on human sperm mitochondria characteristics and their physiological and pathological conditions, paying special attention to improvements in assistant reproductive technology and available treatments to ameliorate male infertility.

Conclusion Although mitochondrial variants associated with male infertility have potential clinical use, research is limited. Further understanding is needed to determine how these characteristics lead to adverse pregnancy outcomes and affect male fertility potential.

Keywords Assisted reproductive technology, In vitro fertilization, Male infertility, Mitochondria, Mitochondrial DNA, Mitochondrial membrane potential, Pregnancy outcomes

[†]Minoo Vahedi Raad and Amir Masoud Firouzabadi are Co-first authors.

*Correspondence:

Ralf Henkel
ralf.henkel@logixxpharma.com; r.henkel@imperial.ac.uk;
rhenkel@uwc.ac.za
Farzaneh Fesahat
farzaneh.fesahat@gmail.com

¹Department of Biology & Anatomical Sciences, Shiraz University of Medical Sciences, Shiraz, Iran

²Reproductive Immunology Research Center, Shahid Sadoughi University of Medical Sciences, Yazd, Iran

³Department of Physiology, School of Medical Sciences, Shahid Sadoughi University of Medical Sciences, Yazd, Iran

⁴Health Reproductive Research Center, Sari Branch, Islamic Azad University, Sari, Iran

⁵LogixX Pharma, Theale, Berkshire, UK

⁶Department of Metabolism, Digestion and Reproduction, Imperial College London, London, UK

⁷Department of Medical Bioscience, University of the Western Cape, Bellville, South Africa



Introduction

Importance of sperm mitochondrial function in male fertility

Globally, about 8–12% of couples suffer from infertility after a year of regular, unprotected intercourse, and 50% of these cases are caused by a male component. Since sperm motility is a crucial requirement for maintaining male fertility, current studies are focusing on the mitochondria as an indicator of normal sperm functionality and fertility [1]. One of the main factors contributing to many cases of “seemingly” idiopathic male infertility is sperm mitochondrial malfunction [2]. Mitochondria are responsible for generating reactive oxygen species (ROS), calcium homeostasis, cell death, and the generation of adenosine triphosphate (ATP) in the process of oxidative phosphorylation [3–5]. It has been suggested that mitochondrial physiology is one of the main factors to maintain chromatin integrity and execute acrosome reactions [6]. Mitochondria contain their own circular DNA (mtDNA) which, due to the absence of DNA-binding proteins such as histones, is about 100 times more susceptible to damage and mutations by high amounts of ROS and matrix-free radicals than nuclear DNA (nDNA) which is protected by histones or, in the case of sperm, by protamines [7]. Additionally, considering that mtDNA replicates quickly and does not have a reliable proof-reading system like the nuclear genome [8], the repair processes for mtDNA are much less effective than those for nDNA. Hence, the mutation rates of mtDNA are reported to be 10 to 17 times higher than for nDNA [9].

Evidence suggests that about 80% of the chromosomal aberrations in humans are of paternal origin [10, 11]. It is widely accepted that in most animals, mtDNA is inherited solely from the mother, with the paternal mitochondria being essentially undetectable after the four-cell stage in a fertilized embryo [12]. This pattern of inheritance is called “maternal inheritance” [13]. However, recent findings cast doubt on this theory as there are indications that extra mtDNA is transferred from fathers to their children [14].

Mitochondrial inheritance

According to a study by Luo and coworkers (2013), maternal factors do not destroy the mitochondria of spermatozoa during early embryonic development [15]. In this specialized polymerase chain reaction (PCR) based study, the authors demonstrated that most motile sperm cells lose their mtDNA during their movement toward the oocyte. However, if sperm with remaining mtDNA entered the oocyte and formed a zygote, this mtDNA was not eliminated but could still be passed on to the progeny and could be detected in the placenta and other extra-embryonic organs [15]. Yet, in order to shield the embryo from the consequences of damage by paternal mtDNA,

sperm mitochondria and mtDNA should be diminished in the early stages of embryonic development [16]. Eker et al. reported that sperm mtDNA is occasionally maintained after whole sperm are injected into the cytoplasm of the oocyte during intra-cytoplasmic sperm injection (ICSI), suggesting that the offspring may actually have their father’s mitochondrial DNA [17]. As a result of intracellular ubiquitination, sperm mitochondria are selectively proteolyzed during early embryonic development [18, 19]. Thus, the proteolytic degradation of sperm mitochondria in the early embryo may be the reason why sperm mitochondria can affect fertilizing potential and early post-fertilization development more than later-stage development [3].

Correlation of mitochondrial dysfunction and infertility

The most accurate measure of mitochondrial activity and the state of the body’s energy supply is the mitochondrial membrane potential (MMP or $\Delta\psi$). Reduced sperm motility and decreased sperm fertilizing capacity are associated with impaired mitochondrial activity as shown by a decrease in sperm MMP [20]. This finding suggests that oxidative phosphorylation, the primary ATP production pathway within mitochondria, is essential for powering sperm motility. However, the study acknowledges the potential role of glycolysis as alternative pathway for energy production. While oxidative phosphorylation likely generates most of the ATP within the mitochondria, glycolysis might play a crucial role in transporting the generated ATP along the sperm’s tail to fuel motility [20]. In addition, some other studies demonstrated an association of sperm mitochondrial dysfunction with recurrent pregnancy loss and the failure of assisted reproductive technology (ART) procedures [21]. Although in vitro fertilization (IVF) is a commonly used method to treat couples with male infertility, its outcome remains unsatisfactory [22]. One of the most critical elements impacting therapeutic outcomes is sperm quality [23]. The latest WHO manual (WHO, 2021) recommends using strict criteria for identifying a normal spermatozoon and gives the following precise definition of a normal spermatozoon: *‘The head should be smooth, regularly contoured and generally oval in shape. acrosomal region comprising 40–70% of the head area without large vacuoles, and not more than two small vacuoles. the post-acrosomal region should not contain any vacuoles. the midpiece should be slender, regular and about the same length as the sperm head. Residual cytoplasm is considered an anomaly only when in excess, i.e. when it exceeds one-third of the sperm head size. principal piece with uniform caliber along its length, must be thinner than the midpiece, and approximately 45 μ m long (about 10 times the head length)’. The success of ICSI, as the most frequently applied method, has led to the use of immature as well as defective sperms*

regardless of subcellular sperm dysfunctions [24]. Failure of oocyte activation is thought to be the most common reason for ICSI failure [25]. Consequently, paternal parameters such as the lack of motile sperm, oocyte activating factors, globozoospermia, sperm DNA damage, and technical issues with sperm injection can all contribute to fertilization failure with ICSI [26]. Different procedures in ART labs have also been suggested to negatively affect sperm mitochondrial health in different ways [27]. Given the conflicting data regarding the impact of sperm mitochondrial dysfunction on pregnancy outcomes and ART success [28, 29], this systematic review aims to critically evaluate existing literature. Our focus is on how a deeper understanding of mitochondrial dysfunction can inform the development of novel ART strategies to improve male infertility treatment. This analysis will encompass a meticulous appraisal of the strengths and limitations within the included studies, facilitating a more nuanced interpretation of the current evidence and pinpointing areas for future research.

Methods

In our literature search, we included the newest original and review scientific literature in the English language, published as full manuscripts (not abstracts) through PubMed, Scopus, and Google Scholar from January 2012 until August 2023. The reporting of this systematic review was guided by the standards of the Preferred Reporting Items for Systematic Review and Meta-Analysis Statement [30]. The following keywords and phrases were used: Male infertility, mitochondria, pregnancy outcomes, assisted reproductive technology, in vitro fertilization, intracytoplasmic sperm injection, sperm quality, mitochondrial DNA, and mitochondrial membrane potential [30].

A total of 2418 articles were identified. After the first screening, 2016 articles were excluded as they were duplicated records or had poor eligibility for irrelevancy of the title or unavailability of the full text. Case reports were also excluded. A total of 402 articles (reviews and original research) were retrieved, of which 282 articles were excluded after full review (Fig. 1). The remaining 120 articles were selected as eligible for the present study: 70 of them were systematic and literature reviews, and 50 original articles.

Results

Based on the studied articles, mitochondrial dysfunction and male infertility are strictly correlated. Mitochondrial impairment can be caused by factors such as oxidative stress, genetics, lifestyle, aging, and damaging ART techniques. Sperm mitochondrial dysfunction alters sperm function and can lead to poor success rates

of ART procedures. Therefore, choosing the proper diagnosis tests is as important as selecting the right treatment option.

Causes of sperm mitochondrial dysfunction

Oxidative stress

Oxidative stress (OS) occurs as a result of an imbalance between elevated ROS levels and the antioxidant capacity in favor of the ROS. Elevated seminal ROS levels may be associated with idiopathic male infertility, thus being a potential screening tool in andrological diagnostics [31]. OS can be correlated with human semen quality, poor fertilization outcomes in ART procedures such as IVF and ICSI, impaired and slower embryo cleavage, poor blastocyst morphology, higher risk of embryonic death, and pre- and post-implantation losses [26, 32].

Sources of ROS in the male reproductive system

In the male reproductive organs, ROS are generated in several compartments such as the seminiferous tubules, epididymis, and in semen by leukocytes as well as to a lesser extent by sperm [33–35].

Any aerobic organism is producing its energy largely in the process of oxidative phosphorylation in the mitochondria. In this process, oxygen accepts electrons, thereby reduced to highly reactive intermediates, and water (H_2O) as the final product is formed. However, during this physiological process, about 1–5% of the consumed oxygen is converted into ROS which are regarded as highly toxic byproducts and are linked to aging and many diseases including infertility [36–39].

Hydrogen peroxide (H_2O_2), superoxide ions ($O_2^{\cdot-}$), and hydroxyl radicals ($\cdot OH$) are the main ROS generated by cells with H_2O_2 mainly involved in redox sensing, signaling, and control [40]. In order to keep the levels of these highly reactive and therefore cytotoxic compounds in balance, antioxidants such as superoxide dismutase, glutathione peroxidase, catalase, glutathione reductase, ascorbate (vitamin C), and α -tocopherol (vitamin E), protect cells from these free radicals, so that the physiologic redox homeostasis is maintained [36]. Even during normal function, the mitochondrial electron transfer chain (ETC) loses electrons, reducing O_2 to $O_2^{\cdot-}$ in a univalent manner. The mitochondrial superoxide dismutase converts 0.1–5% of the oxygen (O_2) that is taken in by the cells into $O_2^{\cdot-}$, which is subsequently quickly changed into H_2O_2 [41, 42]. Physiologically produced ROS at low levels play a crucial role in sperm motility, hyperactivation, transient sperm-oviduct adhesion, controlling sperm capacitation, and acrosome reaction [43, 44]. However, at higher levels ROS may have a negative impact on proteins, lipids, and the DNA, thus significantly harming cell functioning or triggering apoptosis




Fig. 1 Flow chart of study selection according to PRISMA checklist

[40, 45]. Hence, the physiological redox balance, homeostasis, must be maintained.

Other sources of ROS in semen include seminal leukocytes and immature spermatozoa. High quantities of immature spermatozoa exhibiting large amounts of cytoplasm, which is normally phagocytosed by Sertoli cells, result in increased ROS levels. A crucial defensive mechanism against pathogens, the myeloperoxidase system generates ROS from leukocytes in response to inflammation or infection in the male genital tract [46].

Effects of oxidative stress on sperm and mitochondrial function

ROS are produced from physiological metabolic reactions [47]. Even though several organelles including the plasma membrane, endoplasmic reticulum, and peroxisomes produce ROS in cells, mitochondria are the major source of ROS as they also regulate ROS production by non-mitochondrial sources [48–50] and are therefore the cell's main endogenous source of ROS production [51]. When sperm ROS production surpasses the maximum amount of the natural antioxidative protective capacity, processes begin that can harm the sperm mitochondrial membrane and lead to mitochondrial dysfunction [52].

The mitochondrial electron transport chain (ETC) is composed of multiprotein transmembrane complexes (I–IV), generates an electrochemical gradient that enables complex V to synthesize ATP [53]. To function properly and produce ATP during oxidative phosphorylation (OXPHOS), these complexes must be assembled into a specifically configured super complex [41]. Endogenous mitochondrial ROS generation has been connected to the functioning of complexes I and III of the respiratory chain [54–56].

The presence of polyunsaturated fatty acids, which control the fluidity of the membrane and serve as favored targets for oxidative assaults by ROS, makes the sperm plasma membrane highly susceptible to OS-induced damage [57, 58], thus triggering lipid peroxidation and subsequently the formation of highly reactive lipid aldehydes, which create mitochondrial ROS and degrade sperm DNA integrity and competence by covalently attaching to ETC proteins [59]. Although mature sperm were previously considered as not being able to repair their DNA, recent research shows that the male germ cell has a truncated system for detecting and repairing DNA damage involving 8-oxoguanine DNA glycosylase (OGG1). This enzyme was shown to excise 8-hydroxy-2'deoxyguanosine (8OHdG) in the sperm nucleus and mitochondria. Yet, sperm do not possess the downstream enzymatic machinery to complete the repair making sperm more vulnerable to the accumulation of oxidative damage to mitochondrial and nDNA [59]. Additionally,

sperm's cytosolic area is constrained and contains few antioxidants after spermiogenesis [60].

Genetic factors

Mitochondrial genome

Mitochondria have their own genome together with their transcriptional and protein-synthesizing machinery. The mitochondrial genome differs from the nuclear genome in several ways. Mitochondrial DNA is a circular, double-stranded genome that is 16.6 kb long. It contains the genetic material for 22 transfer RNAs, 2 ribosomal RNAs, and 13 polypeptide subunits necessary for the process of oxidative phosphorylation [61]. Within a single cell, there are hundreds to thousands of copies, and introns are not present. Few to no non-coding nucleotides connect its genes, and most of the time termination codons are absent. As highlighted before, due to the rapid replication of mtDNA and the absence of an effective modification mechanism, mtDNA modifications are predicted to be 10 to 17 times greater than that of nDNA (Table 1) [8]. On the other hand, they are produced post-transcriptionally by polyadenylation. Since there are multiple copies of mtDNA, the coexistence of mutant and wild-type molecules with heteroplasmy is possible [62].

The effects of mitochondrial DNA copy number on male fertility

The number of mitochondria differs in somatic cells [66]. Although the number of which is highly dynamic and regulated in a cell-specific manner by mechanisms that are not completely understood, each mitochondrion may contain 1 to 10 copies of the mitochondrial genome [67]. The mtDNA copy number is defined as the proportion of the mitochondrial genome copy numbers to nDNA copies [68]. The number of mtDNA copies in spermatozoa is only about 100. In contrast, oocytes have about 150,000 copies. For healthy sperm function throughout sperm growth and maturation, a reduction in mtDNA copy number is crucial [69].

Many studies have proven the association between increased mtDNA copy number and male infertility, whether it affects sperm parameters and lipid peroxidation in primary fertility or predicts the embryo development process [70–73]. All of these studies support the sensitivity of the mtDNA copy number to oxidative stress and its potential use as a biomarker indicating male fertility potential [70, 74]. It was also shown that infertile men with different sperm parameters have elevated mtDNA content [75]. According to the results of a study by Vahedi Raad et al. on the association between mtDNA and sperm parameters, which was done on two groups of infertile men with normal and abnormal sperm parameters and men who were fertile, this biomarker may be

Table 1 Human nuclear and mitochondrial genome differences

Feature	Nuclear DNA	Mitochondrial DNA	References
Location	Cell nucleus	Mitochondrial matrix	63–65
Size (in bp)	$\sim 3 \times 10^9$	16,569	
Inheritance	Mendelian inheritance for autosomes and X chromosome; Paternal inheritance for Y chromosome	Maternal	
Structure	Double helix, linear, packaged in chromosomes	Double-strand, circular	
Cell division	Cell cycle regulated	Fusion/fission (continuous)	
DNA copies per cell	2 (1 allele from each parent)	~ 10 –50,000	
Total DNA	99.75%	0.25%	
Number of genes encoded	$\sim 20,000$ –30,000 protein coding	13 protein-coding + 24 non-protein coding	
Replication pair	Yes	No	
DNA repair enzymes	Multiple	Polymerase gamma	
Gene density	~ 1 in 40,000 bp	1 in 450 bp	
Introns	Found in almost every gene	Absent	
Histones	Associated with the nuclear DNA	Not associated with the mitochondrial DNA	

utilized to identify and perhaps cure general male infertility [71].

It is unclear how fewer copies of mtDNA would be beneficial for the development of sperms, but because oligozoospermic and asthenozoospermic males have very high levels of sperm mtDNA, scientists hypothesize that the likelihood of ROS-mediated damage is reduced by a decrease in mtDNA content in sperm [76, 77]. Nonetheless, there is no evidence to validate this hypothesis [73]. One hypothesis suggests that increases in the mtDNA copy number may be a healing mechanism by which oxidative damage or mtDNA deletions induce the reproduction of the mitochondrial genome [78]. The mitochondrial-nuclear crosstalk in the male germ cell, which must be kept in balance to drive vital sperm processes including sperm capacitation and sperm-oocyte fusion, becomes dysfunctional when there is excessive ROS production which can then lead to mitochondrial stress because the antioxidant enzyme systems may not be able to regulate the amount of ROS. In turn, this can modify the copy number of mtDNA resulting in mitochondrial DNA mutations, and disrupt the electron transport chain [79].

Atypical gene expression of mtDNA regulators like mitochondria-related mitochondrial transcription factor A (TFAM) or DNA polymerase γ (POLG) may be the cause of the increased mtDNA copy number in defective spermatozoa [80]. Since TFAM expression is normally down-regulated during spermatogenesis, greater levels of mtDNA copy number in aberrant spermatozoa may be related to a physiological aberration in this process [81]. Since the sperm mtDNA copy number was negatively correlated with urine polycyclic aromatic hydrocarbons (PAH) metabolite concentrations in young males, it is suggested that the sperm mtDNA copy number may be responsive to environmental exposures similar to how

the leukocyte mtDNA copy number has been found to be affected by air pollution [70, 82–84]. Therefore, the evaluation of the sperm mtDNA copy number can be used as complementary data for male fertility potential assessment and may help to find the causes of idiopathic male infertility.

Mitochondrial gene variations related to male infertility

There are connections between mitochondrial abnormalities, sperm motility, fertilization rate, and full-term pregnancies because removing mitochondria from sperm and incubating them under oxygen in vitro reduced motility and viability. A significant accumulation of pathogenic mutant mtDNA in the testis results in a reduction in energy generation, ultimately causing meiotic arrest and changes in sperm morphology, thus emphasizing the significance of mitochondrial function in spermatogenesis [85].

The genes for *NADH dehydrogenase 1 (ND1)* and *NADH dehydrogenase 5 (ND5)* are essential components of complex I [86]. Pathogenic variations in the NADH dehydrogenase (ND) genes are anticipated to impact complex I activity, resulting in a shortage in energy generation, which adversely influences sperm motility because complex I is essential for oxidative phosphorylation [87]. The mitochondrial genes ATPase6 and NADH dehydrogenase 4 (ND4) each have two single nucleotide polymorphisms (SNPs) that are related to asthenozoospermia at loci 9055 and 11,719, respectively [88]. Accordingly, the MT-ND4 gene missense mutation (11696G>A) causes the valine residue at position 313 to be replaced with isoleucine, thus changing the protein's secondary structure. This variant has been attributed to decreased sperm motility [89].

An associated protein with cilia and flagella is encoded by the gene Cilia and Flagella Associated Protein 58

(CFAP58), which is primarily expressed in the testis [90]. CFAP58 is a protein that is found throughout the sperm flagella, but it is focused mainly on the middle section, which contains mitochondria. He et al. discovered an association between bi-allelic loss-of-function mutations in CFAP58 and common morphological abnormalities of the sperm flagella. Striking flagellar defects with malformed axonemal and mitochondrial sheaths were seen using transmission electron microscopy. Heat shock protein 60 (HSP60), a protein that forms the mitochondrial sheath, was significantly less abundant in the spermatozoa of males who carried bi-allelic CFAP58 variants, according to immunoblotting and immunofluorescence assays. Thus, CFAP58 deficiency can lead to axoneme and pre-axoneme abnormalities resulting in asthenoteratozoospermia [90].

Mutations and deletions of mtDNA are associated with male infertility [69, 91, 92]. Since mtDNA is present in hundreds to thousands of copies per cell, detecting mutant mtDNA and deletions does not indicate dysfunction as it is generally believed that the mutational load must surpass a threshold value (perhaps exceeding 60% of all mitochondria within a given tissue) for a notable phenotype [93]. The mtDNA 4,977 bp or common deletion, which appears in human mtDNA between nucleotides 8,470 and 13,447, is one of the most well-known mtDNA mutations [94]. This mutation eliminates all or a portion of the genes encoding the five tRNA genes, one complex IV gene, one complex V gene, two complex IV genes, and four complex I genes, all of which are necessary for maintaining normal mitochondrial function [95]. During aging, this deletion accumulates in various tissues and is linked to several illnesses. This has led to its adoption as a marker for mtDNA oxidative damage [96, 97]. The 4,977 bp deletion can be found in 40% of infertile men with mitochondrial dysfunction [98]. While several studies suggested an association between this genetic deletion and sperm parameters [99–101], some other studies could not find this association [72]. Since the frequent deletion of 4,977 bp is connected to aging and is only occasionally present in older males, contradictory additional studies are required for confirmation.

In addition to the typical 4,977 bp deleted mtDNA seen in infertile males, several earlier investigations also found the presence of deletions of 4,866, 4,856, 7,436, and 7,599 bp [92, 102, 103]. Ambulkar and colleagues suggested that the 7,436 bp deletions of mtDNA in spermatozoa may be one of the main causes of dysfunction and immotility of sperm [104]. The origin of these deletions remains debated [105].

Lifestyle factors

Obesity

The prevalence of individuals with obesity is increasing in almost all countries [106]. Obesity and semen quality have both been examined in many studies [107–110]. Researchers found a connection between obesity and reduced mitochondrial activity, sperm progressive motility, and increased DNA fragmentation [111, 112]. Obesity is causing a systemic inflammatory response and is therefore associated with oxidative stress and inflammation [113–116] with elevated seminal ROS levels. The body mass index (BMI) is positively associated with seminal ROS levels [117].

According to an investigation by Binder et al., paternal obesity causes a delay in preimplantation embryo development starting at the second cleavage stage. These authors suggested that this delay is associated with embryonic genome activation. Accordingly, paternal obesity would damage the sperm, possibly at genetic or epigenetic levels [118, 119]. The authors also showed that the MMP in embryos derived from obese males' sperm was significantly decreased as compared to normal sperm [118]. This delay could be due to the sperm from obese men being exposed to high oxidative stress levels as shown by Henkel et al. (not published). Although mitochondrial biogenesis does not occur during the preimplantation phase, mitochondrial DNA is actively transcribed starting at the 2-cell stage to produce respiratory chain components, which would contain proteins encoded by the father's nDNA [120]. According to one study including 344 infertile couples, males with a BMI over 25 kg/m² have a considerably reduced likelihood of clinical pregnancy success than men who are normal weight, however, day 3 embryo quality was unaffected [121]. Another study including 306 couples and 345 normal and overweight males revealed that the normal BMI group had significantly higher sperm activity, semen volume, and total sperm count than the overweight BMI group did. The relationship between obesity and increased oxidative stress, which may result in mitochondrial malfunction, is thought to be the cause of these findings [122]. Therefore, based on evidence from both animal and human models Campbell and co-workers suggested that improving the metabolic health of obese males can improve mitochondrial health and consequently male fertility [123].

Aging

Since it is evident that in recent decades couples tend to have children at older ages [124], it is crucial to understand how aging affects the fertility potential. The importance of maternal age for female fertility potential and reproductive outcome has been proven a long time ago [125]. However, later studies suggest that advanced

paternal age is also associated with declined natural fertility, increased risk of spontaneous pregnancy loss, as well as the health of the offspring [126–131].

Mitochondria are regarded to be crucial to cellular activities that occur throughout the aging process. It is known that mtDNA mutations and mitochondrial dysfunctions increase with age in both animal models and in humans [132]. Although mitochondrial mutation levels increase with age, it is unclear if this increase plays a significant role in the process of aging [132].

The main reason behind mitochondria attribution in the aging process is ROS production [133, 134]. A study conducted by Kang and colleagues indicated that decreased citrate synthase function with age leads to male fertility decline. Citrate synthase is a vital enzyme of the mitochondrial tricarboxylic acid cycle, which directly regulates cellular function [135]. Age-related alterations in fatty acid content and decreased mitochondrial function have been identified in testicular mitochondria, may have an effect on the fluidity and activity of the mitochondrial complex, and are associated with a rise in the formation of superoxide anion, lipid peroxidation, and a decline in the activity of antioxidant enzymes [136–138]. Moreover, the number of Leydig cell mitochondria decreases in aging males [139], thus reducing the ability of Leydig cells to produce testosterone and steroid biosynthesis [140].

Usually, the female reproductive system eliminates defective sperm by natural selection and does not allow their fertilization with eggs. By using intracytoplasmic sperm injection (ICSI), all natural sperm selection barriers are bypassed, thus resulting in an increased risk of passing genetic defects to future generations [141], thereby older men may be more susceptible to this risk, as their ejaculates may contain more damaged sperm. Hence, better strategies should be developed to increase the number of good quality sperm after sperm selection for use in any ART process. Furthermore, in order to fully understand the real extent of the risks of fatherhood at later ages, it is also required to investigate the likelihood of genetic and epigenetic abnormalities in sperm from older men [139].

Lifestyle changes

Mitochondrial function can be negatively or positively affected by many natural dietary polyphenols, depending on the concentration. In addition, Astaxanthin present in seafood can increase MMP and improve sperm mitochondrial function [142]. Carbohydrates and proteins are also nutritional regulators of OS and testosterone levels and are firmly associated with sperm mitochondrial function [143]. Some other studies also suggested the cessation of smoking, especially hookah, reduces oxidative stress and decreases mitochondrial damage [144–146].

Sperm preparation and cryopreservation techniques

In the andrology and embryology laboratory, oxidative stress levels may increase as the procedures such as centrifugation, manipulation of gametes under atmospheric oxygen tension, exposure to light, incubation time, and the type of sperm preparation technique, and/or the type of handling and culture medium used, as well as volatile organic compounds, and culture media additives have been shown to cause additional ROS production [147–153]. The two most frequently employed methods for sperm separation are swim-up and density gradient centrifugation. While the swim-up technique relies on the ability of motile sperm to swim out of the seminal fluid or a pellet of sperm into the supernatant culture medium, density gradient centrifugation selects sperm with normal morphology depending on their specific density [154].

The primary objective of sperm preparation techniques is to separate motile sperm from seminal plasma that contains non-motile, apoptotic, morphologically defective, and dead sperm [155]. When compared with the raw semen, ROS levels may increase throughout the centrifugation stages of certain sperm preparation procedures. In comparison with the swim-up, the chance of picking a spermatozoon with numerous dysfunctions is greater in the post-density gradient [156]. In order to reduce the oxidative stress levels even further, recent sperm selection techniques attempt to mimic the physiological processes in the female reproductive tract by employing microfluidics to reduce sperm oxidative stress and improve DNA fragmentation and the number of top-quality blastocysts [157–160].

The goal of sperm cryopreservation is to preserve spermatozoa of men who need to do so before receiving treatments that affect their ability to conceive such as chemotherapy, radiation, orchiectomy, or vasectomy for future use in ART, and donor insemination programs [161]. Numerous studies report that cryopreservation causes mitochondrial injuries, leading to a decrease in sperm MMP and ATP production [162]. Since mitochondria are the most sensitive sperm organelle to cryopreservation [163], it has been suggested that changes in mitochondrial membrane fluidity caused by cryopreservation increase MMP and induce the release of ROS, which in turn damages the sperm DNA by causing single- or double-strand DNA breaks [164]. The major event during cryopreservation is the disruption of energy supply due to the formation and prolonged opening of mitochondrial permeability transition pores, leading to a sudden collapse of mitochondrial function and reducing sperm motility and the ability to fertilize oocytes [165]. According to recent research on cryopreserved bull sperm, the opening of the mitochondrial permeability transition pore brought about by intracellular Ca^{2+}

increases cause anomalies in mitochondria. This process is also linked to changes in MMP, ATP content, ROS levels, and plasma membrane integrity [166].

During the normal cryopreservation process, ice crystals are formed inside the sperm causing approximately half of the sperm being lost through cell death and lysis. Thus, the quality of the sample after thawing is adversely affected [167]. Techniques to reduce this cellular damage are vitrification and ultra-rapid freezing of the sperm and are therefore widely employed in advanced reproductive medicine [168]. In comparison with conventional cryopreservation protocols, vitrification has been shown to significantly improve sperm motility, recovery of motile sperm, DNA damage, and mitochondrial activity [169].

Effect of sperm mitochondrial dysfunction on fertilization

The effect of mitochondrial defects on sperm motility

Sperm motility is essential for male fertility. In choosing the best course of therapy for an infertile couple, motility is a crucial consideration. A minimum of 5×10^6 sperm must be collected from a semen specimen to recommend intrauterine insemination (IUI); when 2 to 5×10^6 motile sperm can be recovered, IVF is indicated as a therapy; and when samples only yield up to 2×10^6 motile spermatozoa, ICSI is advised [170]. It has been demonstrated that reduced sperm motility can decrease fertilization success and impact the development of early embryos, even if some research suggests that pregnancy or the euploidy rate of the acquired blastocysts in ICSI is not affected. Also, it might be the result of a decreased blastocyst genesis rate or a developmental halt at the cleavage stage (day 3 embryos) [171]. Choosing the most motile sperm is beneficial as sperm motility is positively associated with good sperm DNA integrity [172].

As sperm approach the oocyte, progressive motility, generated by transverse waves moving from proximal to distal along the sperm flagellum, is important. By this mechanism, the sperm is propelled to the oocyte by a hydrodynamic impulse created as a result. Closer to the oocyte, and especially important for the penetration of the oocyte, sperm capacitate and develop hyper-activated motility characterized by an increase in high amplitude, asymmetrical flagellar beat, and increased ATP production [173]. This flagellar movement is driven by ATP, which acts as a molecular motor [174]. The energy is aerobically generated and depends on the oxidative activity of the mitochondria. In light of this, understanding mitochondrial activity may reveal important information about sperm motility [175].

Unfortunately, despite the clear connection between mitochondria and sperm motility, the knowledge has not yet been efficiently applied in clinical practice. Mutations and deletions in the mitochondrial genome impair sperm

motility, resulting in infertility [176] and numerous mitochondrial genome alterations have drawn attention to the significance of mitochondria in male fertility [73]. Kinesin light chain 3 (KLC3), for instance, connects mitochondria, and its failure to bind outer dense fibers results in abnormalities in the mitochondrial sheath, changing progressive motility and causing infertility [177]. Additionally, it has been proposed that KLC3, selenoprotein phospholipid glutathione peroxidase 4 (GPX4), and Golgi-associated PDZ- and coiled-coil motif-containing protein (GOPC) are required to preserve the link between outer dense fibers and mitochondria [178].

The bulk of the defective proteins found in poorly motile sperm were found to be of mitochondrial origin [179]. Some other studies showed that mitochondrial dysfunction is correlated with mitochondrial enzyme activities, oxidative imbalance, and abnormal MMP [175, 180–182]. Furthermore, it was discovered that all of the assessed mitochondrial respiratory complex activities directly and favorably correlate with spermatozoa motility [183], implying that the condition of asthenozoospermia could be caused by changes in how mitochondria operate [20].

The effect of mitophagy impairment on sperm function

To preserve cellular homeostasis under external stress, autophagy has been preserved through the evolution of eukaryotes. Autophagy is classified into three types: macro-, micro-, and chaperone-mediated autophagy [184]. When organelles and proteins are degraded for the production of new macromolecules or as a source of energy, macroautophagy is initiated by a stimulus such as famine and hypoxia [185]. Microautophagy can occur when cytoplasmic material invades the lysosome directly [186]. When heat shock protein 70 (Hsp70) chaperone connects with the lysosome-associated membrane protein type 2a (LAMP-2 A) during chaperone-mediated autophagy, it recognizes proteins with the KFERQ motif causing the attached protein to be translocated into the lysosome and atrophy [187].

Embryogenesis, implantation, and pregnancy maintenance are all significantly influenced by autophagy. Scientists suggest that mitophagy (autophagy of mitochondria) is essential for the differentiation of sperm before fertilization and the removal of paternal mitochondria after fertilization. According to some theories, autophagy controls cell homeostasis during spermatogenesis, changing the proportion of germ cells to Sertoli cells in the testis [188]. For the mitochondria to reorganize properly, extra mitochondria in the residual bodies must be removed by mitophagy during spermatogenesis [189]. In the absence of mitophagy, damaged mitochondria accumulate and cause oxidative stress [190].

When excessive ROS are produced in damaged mitochondria and these mitochondria are not removed by mitophagy, the loss of mitochondrial membrane potential and the release of cytochrome c can trigger apoptosis [191, 192]. The typical indications of cell death, such as cell shrinkage, chromatin condensation and fragmentation, cell vacuolization, and the formation of apoptotic bodies, are not present in spermatozoa. Therefore, the beginning of autophagy in sperm can only be detected with apoptosis cell markers. The function of mitochondria could be negatively impacted by the presence of apoptosis markers in the mid-piece area. As a result, the ability of the male gamete to penetrate the oolemma weakens, which lowers the potency of sperm fertilization [193, 194].

Existing data is consistent with earlier predictions that sperm mitochondria penetrate the mammalian oocyte during fertilization and undergo active degradation inside the oocyte cytoplasm after fertilization [195]. In the majority of eukaryotic species, paternal mitochondria are destroyed by mitophagy through the autophagy receptors p62 and γ -aminobutyric acid receptor-associated protein (GABARAP) after fertilization [130]. When autophagy is inhibited, heteroplasmy formation and the early embryonic stage inheritance of paternal mitochondrial genes occur [196]. Furthermore, mitochondrial damage can spread throughout the body due to the persistence of paternal mtDNA [132]. Hence, the association between male infertility and mitophagy has significant clinical implications for future diagnosis and therapy (Fig. 2).

Success rate of assisted reproductive technologies

Due to the broad use of ART technologies, especially ICSI, which bypasses all natural selection restrictions, accurate investigation to evaluate the health of the male germ cells is crucial in order to prevent the acute financial burden of repeated ART failure and to reduce emotional and psychological stress [197].

In vitro fertilization

Zhang et al. showed that even in normally looking sperm with poor fertilization rates after IVF, mitochondria might be impaired and dysfunctional [198]. Accordingly, sperm mitochondrial modifications can affect ART outcomes because of excess ROS production, mtDNA variations, mutations, deletions, decreased MMP, etc. Wu et al. reported that the sperm mtDNA copy number and mtDNA deletions are related to reduced probabilities of fertilization, high-quality Day 3 embryos, and transfer-quality Day 5 embryos in an ART environment [71]. Conceivably, important characteristics including smoking status and procedure type (IVF vs. ICSI) did not affect the outcomes [71]. Another study conducted by Yang and

colleagues showed that paternal obesity results in lower sperm mitochondrial activity and poor IVF success rates [122].

Liu and colleagues reported that sperm mitochondrial ROS in cleavage-stage embryos have conflicting impacts depending on whether the female patient had normal ovulation. A high percentage of sperm producing a sufficient amount of ROS resulted in good development rates of day-3 embryos in anovulation patients, even though enhanced sperm mitochondrial ROS hindered embryo development in patients with normal ovulation [29]. This could be due to the fact that a small, physiological amount of ROS is necessary to trigger sperm capacitation, and acrosome reaction which are essential for embryo development [199–201]. Yet, further studies to investigate these mechanisms are necessary.

Intracytoplasmic sperm injection

Eker et al. discovered that male offspring after ICSI had the same SNPs in mitochondrial genes such as *cytochrome c oxidase subunit I (COX1)*, *ND1*, *NADH dehydrogenase 4 (ND4)*, and *ND5* as their fathers. In some cases, the degree of similarity reached up to 99% of the paternal mtDNA [17]. Tiegs et al. concluded that the sperm mitochondrial DNA copy number is not associated with fertilization, blastulation, blastocyst euploidy, and live birth rates after IVF and ICSI procedures [202]. In contrast, Wu and colleagues demonstrated that higher sperm mtDNA copy number and mtDNA deletions showed significantly lower odds of fertilization in ICSI treatment [71]. In another study conducted by Sharbatoghli et al., no significant correlation was observed between DNA fragmentation, apoptosis, and dysfunction of sperm MMP and ICSI outcomes [203]. According to research by Ghalei and colleagues, large concentrations of internal H_2O_2 and $O_2^{\cdot-}$ in ICSI patients impact negatively the potential of the mitochondrial membrane. Nevertheless, only H_2O_2 might prevent pronuclear formation [204]. Such contradictory results highlight the urgent need for more detailed studies in this field (Fig. 2).

A study by Torra-Massana and colleagues demonstrated that five mitochondrial proteins have a higher abundance in patients with ICSI failure (dihydrolypoyl transacetylase (DLAT), D subunit of mitochondrial ATP synthase (ATP5H), solute carrier family 25 member 3 (SLC25A3), solute carrier family 25 member 6 (SLC25A6), and fumarate hydratase (FH)). All these proteins are directly linked to the oxidative phosphorylation network [205]. Abnormal mitochondrial function may be linked to reduced proteasome system activity and vice versa. The authors suggested that sperm proteasomal activity is a factor that affects sperm fertilization ability in ICSI.




Fig. 2 Some effects of sperm mitochondrial dysfunction on Fertilization
 GPX4: Selenoprotein phospholipid glutathione peroxidase 4, GOPC: Golgi-associated PDZ- and coiled-coil motif-containing protein, KLC: Kinesin light chain 3, ROS: Reactive oxygen species

Diagnostic tests and treatment of sperm mitochondrial dysfunction

Diagnostic tests

Genetic differences and abnormalities in the mitochondrial genome can be detected by a variety of PCR methods. Furthermore, determination of the sperm energy metabolism of ATP levels is necessary. Measurement of ATP and ADP levels can be achieved using either high-performance liquid chromatography or commercial kits. Mitochondrial calcium levels can also be measured

because Ca^{2+} is considered to be the primary regulator of oxidative phosphorylation. The evaluation of sperm calcium can be performed by fluorescent probes such as Quin-2 AM, fluo-3/AM, and indo-1 AM [206]. Other diagnostic tests are described below.

Mitochondrial activity index test

Nitro-blue tetrazolium is a useful indicator for testing the mitochondrial redox enzyme apparatus. In this test, the water-soluble, yellowish nitro-blue tetrazolium is

converted into an insoluble blue formazan pigment in and around the mid-piece. In this method, sperm with normal mitochondrial activity shows a stained mid-piece, whereas poor staining is a marker of non-motile or poor mitochondrial activity sperm [207]. Some other indirect semi-quantitative tests in which the quantity of color generated would be indirect evidence of mitochondrial activity were also developed. The test was found to have a significant correlation with the sperm motility parameters [208].

Assessment of the mitochondrial membrane potential

Mitochondrial energy storage involves the proton gradients created by the inner mitochondrial membrane and electric potential across this membrane is causing the electron transport across the membranes for ATP synthesis [209].

Studies have shown a strong positive correlation between MMP and ROS production indicating that mitochondria produce more ROS at a high mitochondrial membrane potential [210, 211]. However, lower sperm motility and reproductive capacity are associated with altered mitochondrial activity, which is indicated by a decreased MMP [20]. The MMP is the most precise measure of mitochondrial energy state and function and is correlated with motility as a precise measure of sperm viability and fertilization capacity [212–215]. Due to the lower MMP in poorly motile sperm, it can be used as a factor for IVF success prediction because spermatozoa with low MMP are less capable of experiencing the acrosome reaction and a high percentage of motile spermatozoa is necessary for successful IVF [182]. Besides motility, sperm with high MMP are morphologically normal and can undergo acrosomal reaction. A high mitochondrial membrane potential does not necessarily mean the success of IVE, but if the potential is low, the process will fail and in this case, ICSI would be a more appropriate course of action [215]. Therefore, the determination of sperm MMP is a better criterion than sperm motility and can be recommended as a routine examination to assess sperm quality during diagnostic work-up for assisted reproduction as a supplement to the conventional semen analysis. To evaluate the MMP and sperm quality, fluorescent dyes like JC-1, MitoSOX Red, MitoTracker Green FM (MT-G), and MitoTracker Red CMH2XRos (MT-R) are utilized.

Measurement of oxygen consumption

This test directly assesses the ability of sperm mitochondria to utilize oxygen for energy production. Techniques like respirometry can measure the amount of oxygen consumed by sperm, indicating mitochondrial function. Ruiz-Pesini et al. showed that the mitochondrial membrane potential and oxygen consumption rate are positively associated with the ATP content, the proportion

of motile sperm, and sperm velocity [216]. For the measurement of the oxygen consumption by human spermatozoa, after incubating the sample at 37 °C for at least 30 min, the assay plates are treated with concanavalin A, polyD-lysine, and Cell-Tak adhesive as coating materials to immobilize sperm cells on the bottom well. Next, to optimize the composition of assay media suitable for measuring mitochondrial respiration in human sperm, the XF Cell Mito Stress Test is performed with sperm cells plated in modified Tyrode's solution [217].

Treatment options for mitochondrial dysfunction

Antioxidant therapy

For the treatment of male infertility due to oxidative stress, the use of antioxidants is widely used [218]. The most often used formulations contain zinc, folic acid, N-acetylcysteine, co-enzyme Q10, vitamins E and C, selenium, carnitines, and pentoxifylline, all of which come in a variety of dosages and different combinations [175]. The quality of sperm is improved by supplementing the diet with unsaturated fatty acids such as ω -3 fatty acids, increasing mitochondrial energetic metabolism and decreasing OS [219]. Research has shown that the Mediterranean diet, which contains a high amount of fruits, vegetables, vitamins, proteins, and unsaturated fats, strongly affects sperm health and performance [220].

Selenium is essential for reproductive functions, including testosterone metabolism. It is also a component of the sperm capsule selenoprotein. The supplementation of infertile men with selenium significantly improved sperm motility. Selenoproteins have also been reported to maintain the integrity of sperm morphology [221]. Sperm antioxidant defenses depend on glutathione, which has also been found to improve sperm motility [222]. Selenium and glutathione are vital for phospholipid-hydroperoxide GSH-Px formation which is an enzyme present in sperm cells. This enzyme becomes a structural protein with over 50% of the mitochondrial capsule in the mid-piece of mature spermatozoa. If either substance is deficient, the mid-piece may become unstable, leading to reduced motility [223].

Vitamin E is a crucial antioxidizing substance that primarily protects cell membranes by scavenging ROS and thereby preventing lipid peroxidation. This vitamin is thought to increase the synthesis of antioxidative enzymes that fight free radicals [223].

Since the conventionally used antioxidants are limited in their applications, a new methodology to formulate antioxidant treatments specifically targeting mitochondria as the source of ROS production has been developed. The lipophilic cation, triphenylphosphonium (TPP), is designed to target mitochondria and if conjugated with commonly used antioxidants, TPP-based antioxidants can reach 1000 times higher mitochondrial

antioxidant concentrations [224]. These TTP-based antioxidants largely accumulate in pathological mitochondria to reduce oxidative stress in an MMP-dependent way [225]. Mitoquinone (MitoQ) is one of the well-known mitochondrial-targeted antioxidants. It has been reported that adding MitoQ to the thawing extender can improve sperm quality by decreasing ROS levels and preventing MMP loss in frozen sperm [226].

Assisted reproductive techniques improvement

Melatonin is an antioxidant that is typically produced in mitochondria and accumulated actively through the mitochondrial melatonin transporter. Mitochondria are both the main synthesis site and principal target for melatonin. Melatonin conserves MMP and maintains mitochondrial activities through rummaging of both ROS and RNS, preventing mitochondrial permeability transition pore opening. Furthermore, it functions as a signaling molecule that boosts the expression of genes involved in stress response and antioxidant enzymes [227]. This has led to the supplementation of cryo media with melatonin to avoid cryoinjuries brought on by mitochondrial abnormalities and OS during cryopreservation of the sperm. Melatonin can reduce the harmful reduction in the mitochondrial membrane potential and prevent the inhibition of mitochondrial oxidative phosphorylation and oxidative stress [228–230].

Myoinositol is the most abundant stereoisomer of inositol, influencing the intracellular Ca^{2+} concentration. Recently, Carlomagno and colleagues demonstrated that incubating sperm of oligoasthenoteratozoospermic (OAT) samples with myoinositol greatly enhanced their motility [231]. The enhancement of the intracellular sperm mitochondrial Ca^{2+} concentration due to their incubation in myoinositol containing medium may result in higher inner MMP in sperm cells, which is regarded as an apoptotic marker [232]. In patients with OAT, Rubino and colleagues reported higher motility and fertilization rate in the group where sperm were treated with myoinositol. Thus, myoinositol incubation of sperm might be used to identify patients who benefit from the incubation with myoinositol to increase motility [3].

Conclusion

Spermatozoa rely on healthy mitochondria for ATP production which is essential for motility and fertilization. Mitochondrial dysfunction can lead to infertility. mtDNA damage and mutations can impact sperm function. An increased mtDNA copy number, antioxidants-ROS imbalance, disrupted mitophagy, obesity, and aging harms mitochondrial function leading to poor sperm motility and DNA integrity. Lifestyle changes like dietary modifications and cessation of smoking may improve sperm health. In assisted reproduction, techniques like

centrifugation and cryopreservation can damage sperm mitochondria. Newer methods like microfluidics aim to minimize this damage. Evaluating sperm mitochondrial function, particularly through MMP assessment, can improve ART success after IVF and ICSI. However, further research is needed to understand the link between sperm mitochondrial dysfunction and ART outcomes, develop better methods to evaluate sperm health before ART, and explore therapeutic strategies targeting sperm mitochondria to improve male fertility. By optimizing sperm mitochondrial health, male fertility and success rates in ART procedures can potentially be improved.

Acknowledgements

The authors would like to thank all those who have contributed to this research.

Author contributions

All authors contributed to conception and design. RH and FF were responsible for overall supervision. MVR and AMF and MTN; Drafted the manuscript, which was revised by RH and FF. All authors read and approved the final manuscript.

Funding

The author declares that no funds, grants, or other support were received during the preparation of this manuscript.

Data availability

No datasets were generated or analysed during the current study.

Declarations

Ethical approval

The Shahid Sadoughi University of Medical Sciences approved this study with approval number: IR.SSU.SPH.REC.1402.194.

Consent for publication

Not applicable

Competing interests

The authors declare no competing interests.

Received: 20 April 2024 / Accepted: 27 June 2024

Published online: 17 July 2024

References

1. Moraes CR, Meyers S. The sperm mitochondrion: organelle of many functions. *Anim Reprod Sci.* 2018;194:71–80.
2. La Vignera S, et al. FSH therapy for idiopathic male infertility: four schemes are better than one. *The Aging Male*; 2019.
3. Rubino P, et al. Improving fertilization rate in ICSI cycles by adding myoinositol to the semen preparation procedures: a prospective, bicentric, randomized trial on sibling oocytes. *J Assist Reprod Genet.* 2015;32:387–94.
4. Firouzabadi AM, Rezvani ME, Zare F, Azizian H, Fesahat F. Possible Impact of Human β -defensin 1 on sperm motility in infertile men with abnormal sperm parameters. *Reprod Biol.* 2024;24:100887.
5. Vertika S, Singh KK, Rajender S. Mitochondria, spermatogenesis, and male infertility—An update. *Mitochondrion.* 2020;54:26–40.
6. Durairajanayagam D, et al. Causes and consequences of sperm mitochondrial dysfunction. *Andrologia.* 2021;53(1):e13666.
7. Piomboni P, et al. The role of mitochondria in energy production for human sperm motility. *Int J Androl.* 2012;35(2):109–24.
8. Pandey R, et al. Association of mitochondrial deoxyribonucleic acid mutation with polymorphism in CYP2E1 gene in oral carcinogenesis. *J oral Biology Craniofac Res.* 2012;2(1):4–9.

9. Stewart JB, Chinnery PF. The dynamics of mitochondrial DNA heteroplasmy: implications for human health and disease. *Nat Rev Genet*. 2015;16(9):530–42.
10. González-Marín C, Gosálvez J, Roy R. Types, causes, detection and repair of DNA fragmentation in animal and human sperm cells. *Int J Mol Sci*. 2012;13(11):14026–52.
11. Fesahat F, Firozabadi AM, Zare-Zardini H, Imani M. Roles of different β -defensins in the human reproductive system: a review study. *Am J Men's Health*. 2023;17(3). <https://doi.org/10.1177/15579883231182673>.
12. Cummins J. Fertilization and elimination of the paternal mitochondrial genome. *Hum Reprod*. 2000;15(suppl2):92–101.
13. Sato M, Sato K. Maternal inheritance of mitochondrial DNA by diverse mechanisms to eliminate paternal mitochondrial DNA. (BBA)-Molecular Cell Res. 2013;1833(8):1979–84. *Biochimica et Biophysica Acta*.
14. Luo S, et al. Biparental inheritance of mitochondrial DNA in humans. *Proc Natl Acad Sci*. 2018;115(51):13039–44.
15. Luo S-M, et al. Unique insights into maternal mitochondrial inheritance in mice. *Proc Natl Acad Sci*. 2013;110(32):13038–43.
16. Ford W. Regulation of sperm function by reactive oxygen species. *Hum Reprod Update*. 2004;10(5):387–99.
17. Eker C, et al. Investigation of human paternal mitochondrial DNA transmission in ART babies whose fathers with male infertility. *Eur J Obstet Gynecol Reproductive Biology*. 2019;236:183–92.
18. Ramalho-Santos J. A sperm's tail: the importance of getting it right. *Hum Reprod*. 2011;26(9):2590–1.
19. Nishimura, Y., Yoshinari T, Naruse K, Yamada T, Sumi K, Mitani H, et al. Active digestion of sperm mitochondrial DNA in single living sperm revealed by optical tweezers. *Proc Natl Acad Sci*. 2006;103(5):1382–7.
20. Paoli D, et al. Mitochondrial membrane potential profile and its correlation with increasing sperm motility. *Fertil Steril*. 2011;95(7):2315–9.
21. Imam SN, et al. Idiopathic recurrent pregnancy loss: role of paternal factors; a pilot study. *J Reprod Infertility*. 2011;12(4):267.
22. Sunderam S, et al. Assisted reproductive technology surveillance—United States, 2016. *MMWR Surveillance Summaries*. 2019;68(4):1.
23. Jedrzejczak P, et al. Prediction of spontaneous conception based on semen parameters. *Int J Androl*. 2008;31(5):499–507.
24. Talwar P, Hayatnagar S. Sperm function test. *J Hum Reproductive Sci*. 2015;8(2):61.
25. Meerschaut FV, et al. Assisted oocyte activation following ICSI fertilization failure. *Reprod Biomed Online*. 2014;28(5):560–71.
26. Kuroda S, et al. Early and late paternal effects of reactive oxygen species in semen on embryo development after intracytoplasmic sperm injection. *Syst Biology Reproductive Med*. 2020;66(2):122–8.
27. Ghasemzadeh J, Talebi AR, Khalili MA, Fesahat F, Halvaei I, Nabi A, et al. Sperm parameters, protamine deficiency, and apoptosis in total globozoospermia. *Iran J Reprod Med*. 2015;13(8):495–502.
28. Benkhalifa M, et al. Mitochondria: participation to infertility as source of energy and cause of senescence. *Int J Biochem Cell Biol*. 2014;55:60–4.
29. Liu J, et al. Double-edged sword: effects of human sperm reactive oxygen species on embryo development in IVF cycles. *Reproductive Biology Endocrinol*. 2023;21(1):1–12.
30. Moher D, et al. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Ann Intern Med*. 2009;151(4):264–9.
31. Agarwal A, et al. Effect of oxidative stress on male reproduction. *World J Men's Health*. 2014;32(1):1–17.
32. Tesarik J. Paternal effects on cell division in the preimplantation embryo. *Reprod Biomed Online*. 2005;10(3):370–5.
33. Sharma RK, Agarwal A. Role of reactive oxygen species in male infertility. *Urology*. 1996;48(6):835–50.
34. Suede SH, Malik A, Sapra A. Histology, spermatogenesis. 2020.
35. Wang A, et al. Generation of reactive oxygen species by leukocytes and sperm following exposure to urogenital tract infection. *Arch Androl*. 1997;39(1):11–7.
36. de Kretser DM, et al. Spermatogenesis *Hum Reprod*. 1998;13(suppl1):1–8.
37. Chrousos GP. Stress and disorders of the stress system. *Nat Reviews Endocrinol*. 2009;5(7):374–81.
38. Lloret A, et al. Reductive stress: a new concept in Alzheimer's disease. *Curr Alzheimer Res*. 2016;13(2):206–11.
39. Henkel R, Sandhu IS, Agarwal A. The excessive use of antioxidant therapy: a possible cause of male infertility? *Andrologia*. 2019;51(1):e13162.
40. Sies H, Berndt C, Jones DP. Oxidative stress. *Annu Rev Biochem*. 2017;86:715–48.
41. Zhao RZ, et al. Mitochondrial electron transport chain, ROS generation and uncoupling. *Int J Mol Med*. 2019;44(1):3–15.
42. Cadenas E, Davies KJ. Mitochondrial free radical generation, oxidative stress, and aging. *Free Radic Biol Med*. 2000;29(3–4):222–30.
43. de Lamirande E, O'Flaherty C. Sperm activation: role of reactive oxygen species and kinases. *Biochim et Biophys Acta (BBA)-Proteins Proteom*. 2008;1784(1):106–15.
44. Aitken RJ. Reactive oxygen species as mediators of sperm capacitation and pathological damage. *Mol Reprod Dev*. 2017;84(10):1039–52.
45. Im Yun J, et al. Effects of combined antioxidant supplementation on human sperm motility and morphology during sperm manipulation in vitro. *Fertil Steril*. 2013;100(2):373–8.
46. Koppers AJ, Garg ML, Aitken RJ. Stimulation of mitochondrial reactive oxygen species production by unesterified, unsaturated fatty acids in defective human spermatozoa. *Free Radic Biol Med*. 2010;48(1):112–9.
47. Shadel GS, Horvath TL. Mitochondrial ROS signaling in organismal homeostasis. *Cell*. 2015;163(3):560–9.
48. Plotnikov EY, Zorov DB. Pros and cons of use of mitochondria-targeted antioxidants. 2019, MDPI. p. 316.
49. Collin F. Chemical basis of reactive oxygen species reactivity and involvement in neurodegenerative diseases. *Int J Mol Sci*. 2019;20(10):2407.
50. Zinovkin RA, Zamyatnin AA. Mitochondria-targeted drugs. *Curr Mol Pharmacol*. 2019;12(3):202–14.
51. Bisht S, Dada R. Oxidative stress: major executioner in disease pathology, role in sperm DNA damage and preventive strategies. *Front Biosci (Schol Ed)*. 2017;9(3):420–47.
52. Agarwal A, et al. Characterizing semen parameters and their association with reactive oxygen species in infertile men. *Reproductive Biology Endocrinol*. 2014;12(1):1–9.
53. Guo R, et al. Architecture of human mitochondrial respiratory megacomplex I2III2IV2. *Cell*. 2017;170(6):1247–57. e12.
54. Musset B, et al. NOX5 in human spermatozoa: expression, function, and regulation. *J Biol Chem*. 2012;287(12):9376–88.
55. Koppers AJ, et al. Phosphoinositide 3-kinase signalling pathway involvement in a truncated apoptotic cascade associated with motility loss and oxidative DNA damage in human spermatozoa. *Biochem J*. 2011;436(3):687–98.
56. Thanan R, et al. Oxidative stress and its significant roles in neurodegenerative diseases and cancer. *Int J Mol Sci*. 2014;16(1):193–217.
57. Sikka SC, Rajasekaran M, Hellstrom W. Role of oxidative stress and antioxidants in male infertility. *J Androl*. 1995;16(6):464–8.
58. Henkel RR. Leukocytes and oxidative stress: dilemma for sperm function and male fertility. *Asian J Androl*. 2011;13(1):43.
59. Aitken RJ, Drevet JR. The importance of oxidative stress in determining the functionality of mammalian spermatozoa: a two-edged sword. *Antioxidants*. 2020;9(2):111.
60. Smith TB, et al. The presence of a truncated base excision repair pathway in human spermatozoa that is mediated by OGG1. *J Cell Sci*. 2013;126(6):1488–97.
61. Anderson S, et al. Sequence and organization of the human mitochondrial genome. *Nature*. 1981;290(5806):457–65.
62. Greaves LC, et al. Mitochondrial DNA and disease. *J Pathol*. 2012;226(2):274–86.
63. Larsen NB, Rasmussen M, Rasmussen LJ. Nuclear and mitochondrial DNA repair: similar pathways? *Mitochondrion*. 2005;5(2):89–108. <https://doi.org/10.1016/j.mito.2005.02.002>.
64. Durairajanayagam D, Singh D, Agarwal A, Henkel R. Causes and consequences of sperm mitochondrial dysfunction. *Andrologia*. 2021;53(1):e13666. <https://doi.org/10.1111/and.13666>.
65. Mposhi A, Van der Wijst MG, Faber KN, Rots MG. Regulation of mitochondrial gene expression the epigenetic enigma. *Front Biosci*. 2017;22(7):1099–13. <https://doi.org/10.2741/4535>.
66. Gianotti TF, et al. Mitochondrial DNA copy number is modulated by genetic variation in the signal transducer and activator of transcription 3 (STAT3). *Metabolism*. 2011;60(8):p1142–1149.
67. Robin ED, Wong R. Mitochondrial DNA molecules and virtual number of mitochondria per cell in mammalian cells. *J Cell Physiol*. 1988;136(3):507–13.
68. Montier LLC, Deng JJ, Bai Y. Number matters: control of mammalian mitochondrial DNA copy number. *J Genet Genomics*. 2009;36(3):125–31.
69. Wai T, et al. The role of mitochondrial DNA copy number in mammalian fertility. *Biol Reprod*. 2010;83(1):52–62.

70. Huffman AM, et al. Associations of urinary phthalate metabolites and lipid peroxidation with sperm mitochondrial DNA copy number and deletions. *Environ Res.* 2018;163:10–5.
71. Wu H, et al. Associations of sperm mitochondrial DNA copy number and deletion rate with fertilization and embryo development in a clinical setting. *Hum Reprod.* 2019;34(1):163–70.
72. Raad MV, et al. Altered methyltransferase gene expression, mitochondrial copy number and 4977-bp common deletion in subfertile men with variable sperm parameters. *Andrologia.* 2022;54(10):e14531.
73. Rajender S, Rahul P, Mahdi AA. Mitochondria, spermatogenesis and male infertility. *Mitochondrion.* 2010;10(5):419–28.
74. Rosati AJ, et al. Sperm mitochondrial DNA biomarkers and couple fecundity. *Hum Reprod.* 2020;35(11):2619–25.
75. Zhang G, et al. Mitochondrial biomarkers reflect semen quality: results from the MARCHS study in Chongqing, China. *PLoS ONE.* 2016;11(12):e0168823.
76. Tremellen K. Oxidative stress and male infertility—a clinical perspective. *Hum Reprod Update.* 2008;14(3):243–58.
77. Tian M, et al. Association of DNA methylation and mitochondrial DNA copy number with human semen quality. *Biol Reprod.* 2014;91(4):1–8.
78. Guha M, et al. HnRNPA2 is a novel histone acetyltransferase that mediates mitochondrial stress-induced nuclear gene expression. *Cell Discovery.* 2016;2(1):1–16.
79. Chianese R, Pierantoni R. Mitochondrial reactive oxygen species (ROS) production alters sperm quality. *Antioxidants.* 2021;10(1):92.
80. Amaral A, Ramalho-Santos J, St JC, John. The expression of polymerase gamma and mitochondrial transcription factor A and the regulation of mitochondrial DNA content in mature human sperm. *Hum Reprod.* 2007;22(6):1585–96.
81. Rantanen A, et al. Downregulation of Tfam and mtDNA copy number during mammalian spermatogenesis. *Mamm Genome.* 2001;12:787–92.
82. Ling X, et al. Polycyclic aromatic hydrocarbons exposure decreased sperm mitochondrial DNA copy number: a cross-sectional study (MARHCS) in Chongqing, China. *Environ Pollut.* 2017;220:680–7.
83. Hou L, et al. Airborne particulate matter and mitochondrial damage: a cross-sectional study. *Environ Health.* 2010;9:1–9.
84. Zhong J, et al. Traffic-related air pollution, blood pressure, and adaptive response of mitochondrial abundance. *Circulation.* 2016;133(4):378–87.
85. Tafuri S et al. Reactive oxygen species (ROS) and male fertility. *New Discoveries Embryol.* 2015(2): p. 19–40.
86. Mimaki M, et al. Understanding mitochondrial complex I assembly in health and disease. *Biochim et Biophys Acta (BBA)-Bioenergetics.* 2012;1817(6):851–62.
87. Al Smadi MA, et al. Impact of mitochondrial genetic variants in ND1, ND2, ND5, and ND6 genes on sperm motility and intracytoplasmic sperm injection (ICSI) outcomes. *Reproductive Sci.* 2021;28:1540–55.
88. Holyoake A, et al. High incidence of single nucleotide substitutions in the mitochondrial genome is associated with poor semen parameters in men. *Int J Androl.* 2001;24(3):175–82.
89. Ji J, et al. Mitochondrial DNA sequencing and large-scale genotyping identifies MT-ND4 gene mutation m. 11696G > A associated with idiopathic oligoasthenospermia. *Oncotarget.* 2017;8(32):52975.
90. He X, et al. Bi-allelic loss-of-function variants in CFAP58 cause flagellar axoneme and mitochondrial sheath defects and asthenoteratozoospermia in humans and mice. *Am J Hum Genet.* 2020;107(3):514–26.
91. May-Panloup P, et al. Increased sperm mitochondrial DNA content in male infertility. *Hum Reprod.* 2003;18(3):550–6.
92. Talebi E, Karimian M, Nikzad H. Association of sperm mitochondrial DNA deletions with male infertility in an Iranian population. *Mitochondrial DNA Part A.* 2018;29(4):615–23.
93. Rossignol R, et al. Mitochondrial threshold effects. *Biochem J.* 2003;370(3):751–62.
94. Cortopassi GA, Arnheim N. Detection of a specific mitochondrial DNA deletion in tissues of older humans. *Nucleic Acids Res.* 1990;18(23):6927–33.
95. Nie H, et al. Mitochondrial common deletion, a potential biomarker for cancer occurrence, is selected against in cancer background: a meta-analysis of 38 studies. *PLoS ONE.* 2013;8(7):e67953.
96. Shen L, et al. Evaluating mitochondrial DNA in cancer occurrence and development. *Ann NY Acad Sci.* 2010;1201(1):26–33.
97. Meissner C, et al. The 4977 bp deletion of mitochondrial DNA in human skeletal muscle, heart and different areas of the brain: a useful biomarker or more? *Exp Gerontol.* 2008;43(7):645–52.
98. Shamsi MB, et al. Mitochondrial DNA mutations in etiopathogenesis of male infertility. *Indian J Urology: IJU: J Urol Soc India.* 2008;24(2):150.
99. Al Zoubi MS, et al. 4,977-bp human mitochondrial DNA deletion is associated with asthenozoospermic infertility in Jordan. *Andrologia.* 2020;52(1):e13379.
100. Salehi MH, et al. Low sperm motility due to mitochondrial DNA multiple deletions. *J Chin Clin Med.* 2006;9(1):1–10.
101. Wei Y-H, Kao S-H. Mitochondrial DNA mutation and depletion are associated with decline of fertility and motility of human sperm. Volume 39. *ZOOLOGICAL STUDIES-TAIPEI*; 2000, pp. 1–12. 1.
102. Gholinezhad M, et al. Comparison of large-scale deletions of the sperm mitochondrial DNA in normozoospermic and asthenoteratozoospermic men. *J Cell Biochem.* 2019;120(2):1958–68.
103. Abasalt HC, Gholamali JS, Maryam GC. Lipid peroxidation and large-scale deletions of mitochondrial DNA in asthenoteratozoospermic patients. 2013.
104. Ambulkar PS, et al. Large scale 7436-bp deletions in human sperm mitochondrial DNA with spermatozoa dysfunction and male infertility. *J Clin Diagn Research: JCDR.* 2016;10(11):GC09.
105. Bogueuet M, et al. Mitochondrial DNA content reduction in the most fertile spermatozoa is accompanied by increased mitochondrial DNA rearrangement. *Hum Reprod.* 2022;37(4):669–79.
106. Collaboration NRF. Trends in adult body-mass index in 200 countries from 1975 to 2014: a pooled analysis of 1698 population-based measurement studies with 19.2 million participants. *Lancet.* 2016;387(10026):1377–96.
107. Attaman JA, et al. Dietary fat and semen quality among men attending a fertility clinic. *Hum Reprod.* 2012;27(5):1466–74.
108. Mitchell M, Bakos HW, Lane M. Paternal diet-induced obesity impairs embryo development and implantation in the mouse. *Fertil Steril.* 2011;95(4):1349–53.
109. Tsao C-W, et al. Exploration of the association between obesity and semen quality in a 7630 male population. *PLoS ONE.* 2015;10(3):e0119458.
110. Wang S, et al. Does obesity based on body mass index affect semen quality?—a meta-analysis and systematic review from the general population rather than the infertile population. *Andrologia.* 2021;53(7):e14099.
111. Fariello RM, et al. Association between obesity and alteration of sperm DNA integrity and mitochondrial activity. *BJU Int.* 2012;110(6):863–7.
112. Oliveira J, et al. Association between body mass index and sperm quality and sperm DNA integrity. A large population study. *Andrologia.* 2018;50(3):e12889.
113. Huang C-J, et al. Obesity-related oxidative stress: the impact of physical activity and diet manipulation. *Sports medicine-open.* 2015;1:1–12.
114. Leisegang K. Oxidative stress in men with obesity, metabolic syndrome and type 2 diabetes mellitus: mechanisms and management of reproductive dysfunction. *Oxidative stress and toxicity in Reproductive Biology and Medicine: a comprehensive update on male infertility-volume one.* Springer; 2022. pp. 237–56.
115. Leisegang K, Henkel R, Agarwal A. Obesity and metabolic syndrome associated with systemic inflammation and the impact on the male reproductive system. *Am J Reprod Immunol.* 2019;82(5):e13178.
116. Bhattacharya K, et al. Obesity, systemic inflammation and male infertility. *Chem Biology Lett.* 2020;7(2):92–8.
117. Tunc O, Bakos H, Tremellen K. Impact of body mass index on seminal oxidative stress. *Andrologia.* 2011;43(2):121–8.
118. Sousa AP, et al. Not all sperm are equal: functional mitochondria characterize a subpopulation of human sperm with better fertilization potential. *PLoS ONE.* 2011;6(3):e18112.
119. Binder NK, Mitchell M, Gardner DK. Parental diet-induced obesity leads to retarded early mouse embryo development and altered carbohydrate utilisation by the blastocyst. *Reprod Fertility Dev.* 2012;24(6):804–12.
120. Binder NK, Hannan NJ, Gardner DK. Paternal diet-induced obesity retards early mouse embryo development, mitochondrial activity and pregnancy health. *PLoS ONE.* 2012;7(12):e52304.
121. Merhi ZO, et al. Male adiposity impairs clinical pregnancy rate by in vitro fertilization without affecting day 3 embryo quality. *Obesity.* 2013;21(8):1608–12.
122. Yang Q, et al. Effect of paternal overweight or obesity on IVF treatment outcomes and the possible mechanisms involved. *Sci Rep.* 2016;6(1):29787.
123. Campbell JM, et al. Paternal obesity negatively affects male fertility and assisted reproduction outcomes: a systematic review and meta-analysis. *Reprod Biomed Online.* 2015;31(5):593–604.
124. Toriello HV, et al. Statement on guidance for genetic counseling in advanced paternal age. *Genet Sci.* 2008;10(6):457–60.

125. Mohammadzadeh M, et al. Influential effect of age on oocyte morphometry, fertilization rate and embryo development following IVF in mice. *Middle East Fertility Soc J*. 2018;23(2):117–20.
126. Aitken RJ, et al. Oxidative stress and male reproductive health. *Asian J Androl*. 2014;16(1):31.
127. Johnson SL, Gemmell NJ. Are old males still good males and can females tell the difference? Do hidden advantages of mating with old males offset costs related to fertility, or are we missing something else? *BioEssays*. 2012;34(7):609–19.
128. Kong A, et al. Rate of de novo mutations and the importance of father's age to disease risk. *Nature*. 2012;488(7412):471–5.
129. Dain L, Auslander R, Dirnfeld M. The effect of paternal age on assisted reproduction outcome. *Fertil Steril*. 2011;95(1):1–8.
130. Kovac JR, et al. The effects of advanced paternal age on fertility. *Asian J Androl*. 2013;15(6):723.
131. Chapuis A, et al. Sperm quality and paternal age: effect on blastocyst formation and pregnancy rates. *Basic Clin Androl*. 2017;27(1):1–9.
132. Sun N, Youle RJ, Finkel T. The mitochondrial basis of aging. *Mol Cell*. 2016;61(5):654–66.
133. Guthrie H, Welch G. Effects of reactive oxygen species on sperm function. *Theriogenology*. 2012;78(8):1700–8.
134. Lima T, et al. Pleiotropic effects of mitochondria in aging. *Nat Aging*. 2022;2(3):199–213.
135. Kang W, et al. Extra-mitochondrial citrate synthase initiates calcium oscillation and suppresses age-dependent sperm dysfunction. *Lab Invest*. 2020;100(4):583–95.
136. Vázquez-Memije ME, et al. Respiratory chain complexes and membrane fatty acids composition in rat testis mitochondria throughout development and ageing. *Exp Gerontol*. 2005;40(6):482–90.
137. Vázquez-Memije ME, et al. Analysis of age-associated changes in mitochondrial free radical generation by rat testis. *Mol Cell Biochem*. 2008;307:23–30.
138. Sahoo D, Roy A, Chainy G. Rat testicular mitochondrial antioxidant defence system and its modulation by aging. *Acta Biol Hung*. 2008;59(4):413–24.
139. Amaral S, Amaral A, Ramalho-Santos J. Aging and male reproductive function: a mitochondrial perspective. *Front Biosci (Schol Ed)*. 2013;5(1):181–97.
140. Chen H, et al. Age-related increase in mitochondrial superoxide generation in the testosterone-producing cells of Brown Norway rat testes: relationship to reduced steroidogenic function? *Exp Gerontol*. 2001;36(8):1361–73.
141. Hamdan M, et al. The impact of endometrioma on IVF/ICSI outcomes: a systematic review and meta-analysis. *Hum Reprod Update*. 2015;21(6):809–25.
142. Wolf AM, et al. Astaxanthin protects mitochondrial redox state and functional integrity against oxidative stress. *J Nutr Biochem*. 2010;21(5):381–9.
143. Ferramosca A, Zara V. Diet and male fertility: the impact of nutrients and antioxidants on sperm energetic metabolism. *Int J Mol Sci*. 2022;23(5):2542.
144. Tofghi Niaki M, et al. Possible harmful effects of Smoking Hookah on sperm DNA fragmentation index and protamine genes expression in normozoospermic men. *Subst Abuse: Res Treat*. 2023;17:11782218221144547.
145. Qutub J et al. Smoking and infertility in Saudi Arabian males: a systematic. 2022.
146. Adetona O et al. The adverse health effects of waterpipe smoking in adolescents and young adults: a narrative review. *Tob Induc Dis*, 2021. 19.
147. Anbari F, et al. The quality of sperm preparation medium affects the motility, viability, and DNA integrity of human spermatozoa. *J Hum Reproductive Sci*. 2016;9(4):254.
148. Talevi R, et al. Protective effects of in vitro treatment with zinc, d-aspartate and coenzyme q10 on human sperm motility, lipid peroxidation and DNA fragmentation. *Reproductive Biology Endocrinol*. 2013;11:1–10.
149. Aitken RJ, Clarkson JS. Significance of reactive oxygen species and antioxidants in defining the efficacy of sperm preparation techniques. *J Androl*. 1988;9(6):367–76.
150. Shahar S, et al. Light-mediated activation reveals a key role for protein kinase A and sarcoma protein kinase in the development of sperm hyper-activated motility. *Hum Reprod*. 2011;26(9):2274–82.
151. Muratori M, et al. Sperm selection with density gradient centrifugation and swim up: effect on DNA fragmentation in viable spermatozoa. *Sci Rep*. 2019;9(1):1–12.
152. Selvam MKP, et al. The effect of oxidative and reductive stress on semen parameters and functions of physiologically normal human spermatozoa. *Free Radic Biol Med*. 2020;152:375–85.
153. Agarwal A, et al. Oxidative stress and assisted reproduction: a comprehensive review of its pathophysiological role and strategies for optimizing embryo culture environment. *Antioxidants*. 2022;11(3):477.
154. Pinto S, et al. Sperm selection strategies and their impact on assisted reproductive technology outcomes. *Andrologia*. 2021;53(2):e13725.
155. Raad G, et al. Adverse effects of paternal obesity on the motile spermatozoa quality. *PLoS ONE*. 2019;14(2):e0211837.
156. Raad G, et al. Differential impact of four sperm preparation techniques on sperm motility, morphology, DNA fragmentation, acrosome status, oxidative stress, and mitochondrial activity: a prospective study. *Andrology*. 2021;9:1549–1559.
157. Quinn MM, et al. Microfluidic sorting selects sperm for clinical use with reduced DNA damage compared to density gradient centrifugation with swim-up in split semen samples. *Hum Reprod*. 2018;33(8):1388–93.
158. Keskin M, et al. Does microfluidic sperm sorting affect embryo euploidy rates in couples with high sperm DNA fragmentation? *Reproductive Sci*. 2022;29(6):1801–8.
159. Stone L. Microfluidic sperm selection. *Nat Reviews Urol*. 2023;20(1):7–7.
160. Vasilescu SA, et al. Sperm quality metrics were improved by a biomimetic microfluidic selection platform compared to swim-up methods. *Volume 9. Microsystems & Nanoengineering*; 2023. p. 37. 1.
161. Redig AJ, et al. Incorporating fertility preservation into the care of young oncology patients. *Wiley Online Library*; 2011. pp. 1–10.
162. Merino O, et al. Protective effect of butylated hydroxytoluene on sperm function in human spermatozoa cryopreserved by vitrification technique. *Andrologia*. 2015;47(2):186–93.
163. Pena F, et al. Antioxidant supplementation in vitro improves boar sperm motility and mitochondrial membrane potential after cryopreservation of different fractions of the ejaculate. *Anim Reprod Sci*. 2003;78(1–2):85–98.
164. Said TM, Gaglani A, Agarwal A. Implication of apoptosis in sperm cryoinjury. *Reprod Biomed Online*. 2010;21(4):456–62.
165. Fang Y, Zhang J. Inhibition of mitochondrial permeability transition pore improved quality of frozen-thawed ram sperm. *Cryobiology*. 2018;81:240.
166. Treulen F, et al. Cryopreservation induces mitochondrial permeability transition in a bovine sperm model. *Cryobiology*. 2018;83:65–74.
167. Agha-Rahimi A, et al. Vitrification is not superior to rapid freezing of normozoospermic spermatozoa: effects on sperm parameters, DNA fragmentation and hyaluronan binding. *Reprod Biomed Online*. 2014;28(3):352–8.
168. Satirapod C, et al. Comparison of cryopreserved human sperm from solid surface vitrification and standard vapor freezing method: on motility, morphology, vitality and DNA integrity. *Andrologia*. 2012;44:786–90.
169. Pabón D, et al. A new system of sperm cryopreservation: evaluation of survival, motility, DNA oxidation, and mitochondrial activity. *Andrology*. 2019;7(3):293–301.
170. Dcunha R et al. Current insights and latest updates in sperm motility and associated applications in assisted reproduction. *Reproductive Sci*, 2020: p. 1–19.
171. Loutradi KE, et al. The effects of sperm quality on embryo development after intracytoplasmic sperm injection. *J Assist Reprod Genet*. 2006;23:69–74.
172. Nosrati R, et al. Rapid selection of sperm with high DNA integrity. *Lab Chip*. 2014;14(6):1142–50.
173. Suarez SS. Control of hyperactivation in sperm. *Hum Reprod Update*. 2008;14(6):647–57.
174. Ford W. Glycolysis and sperm motility: does a spoonful of sugar help the flagellum go round? *Hum Reprod Update*. 2006;12(3):269–74.
175. Barbagallo F, et al. Evaluation of sperm mitochondrial function: a key organelle for sperm motility. *J Clin Med*. 2020;9(2):363.
176. Gabriel MS, et al. Influence of microsurgical varicocelelectomy on human sperm mitochondrial DNA copy number: a pilot study. *J Assist Reprod Genet*. 2012;29:759–64.
177. Zhang Y, et al. KLC3 is involved in sperm tail midpiece formation and sperm function. *Dev Biol*. 2012;366(2):101–10.
178. Mi Y, Shi Z, Li J. Spata19 is critical for sperm mitochondrial function and male fertility. *Mol Reprod Dev*. 2015;82(11):907–13.
179. Amaral A, et al. Identification of proteins involved in human sperm motility using high-throughput differential proteomics. *J Proteome Res*. 2014;13(12):5670–84.
180. Ruiz-Pesini E, et al. Correlation of sperm motility with mitochondrial enzymatic activities. *Clin Chem*. 1998;44(8):1616–20.
181. Castellini C, et al. Pathophysiology of mitochondrial dysfunction in human spermatozoa: focus on energetic metabolism, oxidative stress and apoptosis. *Antioxidants*. 2021;10(5):695.
182. Agnihotri SK, et al. Mitochondrial membrane potential (MMP) regulates sperm motility. *Vitro Cell Dev Biology-Animal*. 2016;52:p953–960.

183. Guo H, et al. Relationships between mitochondrial DNA content, mitochondrial activity, and boar sperm motility. *Theriogenology*. 2017;87:276–83.
184. Mizushima N, Komatsu M. Autophagy: Renovation Cells Tissues Cell. 2011;147(4):728–41.
185. Hamasaki M, et al. Autophagosomes form at ER–mitochondria contact sites. *Nature*. 2013;495(7441):389–93.
186. Pohl C, Dikic I. Cellular quality control by the ubiquitin-proteasome system and autophagy. *Science*. 2019;366(6467):818–22.
187. Boya P, Reggiori F, Codogno P. Emerging regulation and functions of autophagy. *Nat Cell Biol*. 2013;15(7):713–20.
188. Aitken RJ, et al. Apoptosis in the germ line. *Reproduction*. 2011;141(2):139.
189. Huang Q, et al. Autophagy core protein ATG5 is required for elongating spermatid development, sperm individualization and normal fertility in male mice. *Autophagy*. 2021;17(7):1753–67.
190. De Gaetano A, et al. Mitophagy and oxidative stress: the role of aging. *Antioxidants*. 2021;10(5):794.
191. Ferraro E, Cecconi F. Autophagic and apoptotic response to stress signals in mammalian cells. *Arch Biochem Biophys*. 2007;462(2):210–9.
192. Yefimova MG, et al. Autophagy is increased in cryptorchid testis resulting in abnormal spermatozoa. *Asian J Androl*. 2019;21(6):570.
193. Talarczyk-Desole J, et al. Sperm midpiece apoptotic markers: impact on fertilization potential in in vitro fertilization and intracytoplasmic sperm injection. *Hum Cell*. 2016;29:67–75.
194. Dehghanpour F, et al. Evaluation of sperm protamine deficiency and apoptosis in infertile men with idiopathic teratozoospermia. *Clin Experimental Reproductive Med*. 2017;44(2):73.
195. Song W-H et al. Regulation of mitochondrial genome inheritance by autophagy and ubiquitin-proteasome system: implications for health, fitness, and fertility. *BioMed research international*. 2014;2014.
196. Al Rawi S, et al. Postfertilization autophagy of sperm organelles prevents paternal mitochondrial DNA transmission. *Science*. 2011;334(6059):1144–7.
197. Morreale M, et al. The impact of stress and psychosocial interventions on assisted reproductive technology outcome. *J Sex Marital Ther*. 2010;37(1):56–69.
198. Zhang X-Y, et al. Melatonin rescues impaired penetration ability of human spermatozoa induced by mitochondrial dysfunction. *Reproduction*. 2019;158(5):465–75.
199. Ufer C, Wang CC. The roles of glutathione peroxidases during embryo development. *Front Mol Neurosci*. 2011;4:11531.
200. Belli M, et al. Oxygen concentration alters mitochondrial structure and function in in vitro fertilized preimplantation mouse embryos. *Hum Reprod*. 2019;34(4):601–11.
201. Deluao JC, et al. OXIDATIVE STRESS AND REPRODUCTIVE FUNCTION: reactive oxygen species in the mammalian pre-implantation embryo. *Reproduction*. 2022;164(6):F95–108.
202. Tiegs AW, et al. Sperm mitochondrial DNA copy number is not a predictor of intracytoplasmic sperm injection (ICSI) cycle outcomes. *Reproductive Sci*. 2020;27(6):1350–6.
203. Sharbatoghli M, et al. Relationship of sperm DNA fragmentation, apoptosis and dysfunction of mitochondrial membrane potential with semen parameters and ART outcome after intracytoplasmic sperm injection. *Arch Gynecol Obstet*. 2012;286:1315–22.
204. Ghaleno L, et al. High level of intracellular sperm oxidative stress negatively influences embryo pronuclear formation after intracytoplasmic sperm injection treatment. *Andrologia*. 2014;46(10):1118–27.
205. Torra-Massana M, et al. Altered mitochondrial function in spermatozoa from patients with repetitive fertilization failure after ICSI revealed by proteomics. *Andrology*. 2021;9(4):1192–204.
206. de Agostini Losano JD, et al. Spermatic mitochondria: role in oxidative homeostasis, sperm function and possible tools for their assessment. *Zygote*. 2018;26(4):251–60.
207. Gosálvez J, Fernández JL, Esteves SC. Response: Nitroblue tetrazolium (NBT) assay. *Reprod Biomed Online*. 2018;36(1):92–3.
208. Nowicka-Bauer K et al. Sperm mitochondrial dysfunction and oxidative stress as possible reasons for isolated asthenozoospermia. *J Physiol Pharmacol*, 2018. 69(3).
209. Marchetti C, et al. Study of mitochondrial membrane potential, reactive oxygen species, DNA fragmentation and cell viability by flow cytometry in human sperm. *Hum Reprod*. 2002;17(5):1257–65.
210. Zorova LD, et al. Mitochondrial membrane potential. *Anal Biochem*. 2018;552:50–9.
211. Suski J et al. Relation between mitochondrial membrane potential and ROS formation. *Mitochondrial bioenergetics: Methods and Protocols*. 2018:357–381.
212. Kasai T, et al. Relationship between sperm mitochondrial membrane potential, sperm motility, and fertility potential. *Asian J Androl*. 2002;4(2):97–103.
213. Marchetti C, et al. Comparison of four fluorochromes for the detection of the inner mitochondrial membrane potential in human spermatozoa and their correlation with sperm motility. *Hum Reprod*. 2004;19(10):2267–76.
214. Espinoza J, et al. Integrity of mitochondrial membrane potential reflects human sperm quality. *Andrologia*. 2009;41(1):51–4.
215. Marchetti P, et al. Influence of mitochondrial membrane potential of spermatozoa on in vitro fertilisation outcome. *Andrologia*. 2012;44(2):136–41.
216. Ruiz-Pesini E, et al. The role of the mitochondrion in sperm function: is there a place for oxidative phosphorylation or is this a purely glycolytic process? *Curr Top Dev Biol*. 2007;77:3–19.
217. Taniguchi H, et al. Establishment of a novel assessment of the quality of human spermatozoa measuring mitochondrial oxygen metabolism. *BMC Res Notes*. 2022;15(1):123.
218. Agarwal A, et al. Impact of antioxidant therapy on natural pregnancy outcomes and semen parameters in infertile men: a systematic review and meta-analysis of randomized controlled trials. *World J Men's Health*. 2023;41(1):14.
219. Ferramosca A, et al. Dietary fatty acids influence sperm quality and function. *Andrology*. 2017;5(3):423–30.
220. Karayiannis D, et al. Association between adherence to the Mediterranean diet and semen quality parameters in male partners of couples attempting fertility. *Hum Reprod*. 2017;32(1):215–22.
221. Qazi IH, et al. Role of selenium and selenoproteins in male reproductive function: a review of past and present evidences. *Antioxidants*. 2019;8(8):268.
222. ET D. Glutathione and hypotaurine in vitro: effects on human sperm motility, DNA integrity and production of reactive oxygen species. *Mutagenesis*. 2000;15:61–8.
223. Moslemi MK, Tavanbakhsh S. Selenium–vitamin E supplementation in infertile men: effects on semen parameters and pregnancy rate. *Int J Gen Med*, 2011: p. 99–104.
224. Zielonka J, et al. Mitochondria-targeted triphenylphosphonium-based compounds: syntheses, mechanisms of action, and therapeutic and diagnostic applications. *Chem Rev*. 2017;117(15):10043–120.
225. Wang JY, et al. Triphenylphosphonium (TPP)-Based antioxidants: a new perspective on antioxidant design. *ChemMedChem*. 2020;15(5):404–10.
226. Kumar P, et al. Unraveling subcellular and ultrastructural changes during vitrification of human spermatozoa: Effect of a mitochondria-targeted antioxidant and a permeable cryoprotectant. *Front Cell Dev Biology*. 2021;9:672862.
227. Tan D-X, et al. Melatonin: a mitochondrial targeting molecule involving mitochondrial protection and dynamics. *Int J Mol Sci*. 2016;17(12):2124.
228. Fang Y, et al. Melatonin inhibits formation of mitochondrial permeability transition pores and improves oxidative phosphorylation of frozen-thawed ram sperm. *Front Endocrinol*. 2020;10:896.
229. Leon J, et al. Melatonin mitigates mitochondrial malfunction. *J Pineal Res*. 2005;38(1):1–9.
230. Govender J, et al. Mitochondrial catastrophe during doxorubicin-induced cardiotoxicity: a review of the protective role of melatonin. *J Pineal Res*. 2014;57(4):367–80.
231. Carlomagno G, et al. Contribution of myo-inositol and melatonin to human reproduction. *Eur J Obstet Gynecol Reproductive Biology*. 2011;159(2):267–72.
232. Condorelli RA, et al. Myo-inositol: does it improve sperm mitochondrial function and sperm motility? *Urology*. 2012;79(6):1290–5.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.