



CrossMark

electronic reprint

Acta Crystallographica Section C

**Crystal Structure
Communications**

ISSN 0108-2701

Editor: **Anthony Linden**

Dichlorido{*N*-[2-(diphenylphosphanyl)benzylidene]-2-methylaniline}palladium(II) acetonitrile monosolvate


William M. Motswainyana, Martin O. Onani, Jeroen Jacobs and Luc Van Meervelt

Acta Cryst. (2013). **C69**, 209–211

Copyright © International Union of Crystallography

Author(s) of this paper may load this reprint on their own web site or institutional repository provided that this cover page is retained. Republication of this article or its storage in electronic databases other than as specified above is not permitted without prior permission in writing from the IUCr.

For further information see <http://journals.iucr.org/services/authorrights.html>



Acta Crystallographica Section C: Crystal Structure Communications specializes in the rapid dissemination of high-quality studies of crystal and molecular structures of interest in fields such as chemistry, biochemistry, mineralogy, pharmacology, physics and materials science. The numerical and text descriptions of each structure are submitted to the journal electronically as a Crystallographic Information File (CIF) and are checked and typeset automatically prior to peer review. The journal is well known for its high standards of structural reliability and presentation. *Section C* publishes approximately 1000 structures per year; readers have access to an archive that includes high-quality structural data for over 10000 compounds.

Crystallography Journals Online is available from journals.iucr.org

Dichlorido{*N*-[2-(diphenylphosphanyl)benzylidene]-2-methylaniline}-palladium(II) acetonitrile monosolvate

William M. Motswainyana,^a Martin O. Onani,^a Jeroen Jacobs^b and Luc Van Meervelt^{b*}

^aUniversity of the Western Cape, Private Bag X17, Bellville 7535, South Africa, and

^bChemistry Department, KU Leuven, Celestijnenlaan 200F, B-3001 Leuven (Heverlee), Belgium

Correspondence e-mail: luc.vanmeervelt@chem.kuleuven.be

Received 21 January 2013

Accepted 23 January 2013

Online 5 February 2013

The title imino-phosphine compound, $[\text{PdCl}_2(\text{C}_{26}\text{H}_{22}\text{NP})]\cdot\text{CH}_3\text{CN}$, was prepared by reaction of *N*-[2-(diphenylphosphanyl)benzylidene]-2-methylaniline with dichlorido(cycloocta-1,5-diene)palladium(II) in dry CH_2Cl_2 . The Pd^{II} cation is coordinated by the P and N atoms of the bidentate chelating ligand and by two chloride anions, generating a distorted square-planar coordination geometry. There is a detectable *trans* influence for the chloride ligands. The methyl group present in this structure has an influence on the crystal packing.


Comment

Imino-phosphine complexes have been studied since the early 1990s (Newkome, 1993; Ghilardi *et al.*, 1992). Their preparation is versatile because a wide range of amines and aldehydes of varying steric bulk are either available commercially or can be easily synthesized (Mogorosi *et al.*, 2011; Nobre & Monteiro, 2009; Pellagati *et al.*, 2005; Newkome, 1993; Ghilardi *et al.*, 1992; Vaughan *et al.*, 2011; Puri *et al.*, 2011; Doherty *et al.*, 2002; Onani *et al.*, 2010; Motswainyana *et al.*, 2012). The soft P atom coordinates strongly to soft metals, while the hard N atom is weakly coordinated and therefore more easily displaced, thus allowing hemilability. The hemilability of these ligands provides a unique reversible protection of coordination sites, an important property of ligands which leads to improved catalytic reactions (Ojwach *et al.*, 2007).

Phosphine-based complexes have been less well explored for their potential to inhibit tumour growth, possibly due to their susceptibility to oxidation, which makes them very difficult to handle in air. In our attempt to prepare new imino-phosphine palladium(II) complexes, which could induce apoptosis in tumour cells, we synthesized and crystallized the title compound, (I).

Compound (I) has proved to possess remarkable anti-proliferative activities against the human breast (MCF-7) and

human colon (HT-29) cancer cell lines compared with the reference drug cisplatin. The recorded cytotoxic activities could be attributed to the higher aqueous solubility of this palladium(II) complex, which makes it easier to dissociate in solution, thereby making the compound bio-available (Gao *et al.*, 2009). The imino-phosphine ligand *N*-[2-(diphenylphosphanyl)benzylidene]-2-methylaniline forms a six-membered ring around the metal centre, which stabilizes the complex during biological investigations, probably resulting in the observed antitumour profiles (Bacchi *et al.*, 2000).



The Pd^{II} cation in (I) (Fig. 1) is coordinated by the P and N atoms of the imino-phosphine ligand and two chloride anions, generating a distorted square-planar coordination geometry around the metal centre. The bond angles around $\text{Pd}1$ (Table 1) describe the observed distorted square-planar geometry, while also indicating some ring strain induced by the chelating nature of the bidentate ligand. The $\text{Pd}1-\text{C}1$ bond lengths (Table 1) are in good agreement with the average $\text{Pd}-\text{Cl}$ bond length of 2.298 (15) Å for known palladium(II) complexes (Allen, 2002; Chiririwa & Muller, 2012). The average $\text{Pd}-\text{N}$ and $\text{Pd}-\text{P}$ bond lengths (Table 1) also compare well with literature values (Chiririwa & Muller, 2012; Motswainyana *et al.*, 2012). There is a detectable *trans* influence for the chloride anions, since the $\text{Pd}1-\text{Cl}2$ bond is significantly longer than $\text{Pd}1-\text{Cl}1$, thus reflecting the stronger *trans* influence of the diphenylphosphanyl group compared to the amine (Doherty *et al.*, 2002). The Pd^{II} cation deviates by 0.0026 (3) Å from the best least-squares plane through atoms P1, N1, Cl1 and Cl2. This plane makes an angle of 46.57 (11)° with the best least-squares plane through the C2–C7 ring. The Pd-containing six-membered ring has a screw-boat confor-




Figure 1

The molecular structure of (I), showing the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level.

mation, with atoms Pd1 and P1 deviating from the plane through the other four atoms.

Despite the presence of several benzene rings, no $\pi\cdots\pi$ interactions are present in (I); only a number of weaker C—H $\cdots\pi$ contacts are observed [C23—H23 \cdots Cg1ⁱ = 2.84 Å and C26—H26A \cdots Cg2ⁱⁱ = 2.93 Å; Cg1 and Cg2 are the centroids of the C2—C7 and C14—C19 rings, respectively; symmetry codes: (i) $-x + \frac{3}{2}, y - \frac{1}{2}, -z + \frac{3}{2}$; (ii) $-x + \frac{3}{2}, y + \frac{1}{2}, -z + \frac{3}{2}$].

The asymmetric unit of (I) contains one acetonitrile solvent molecule, which was essential during the crystallization experiments to obtain good quality crystals. Two acetonitrile molecules occupy relatively large voids of 233 Å³, and there are four such voids in the unit cell. In the packing, the position of the acetonitrile is fixed on one side by a C27—H27C \cdots Cl2 interaction [H27C \cdots Cl2ⁱⁱⁱ = 2.70 Å; symmetry code: (iii) $-x + 1, -y + 2, -z + 1$] and by a weak π -interaction (C27—H27B \cdots Cg2 = 3.29 Å) (Fig. 2). Furthermore, atom N2 interacts with atoms H11, H16 and H22 of neighbouring molecules [N2 \cdots H11^{iv} = 2.76 Å, N2 \cdots H16^v = 2.65 Å and N2 \cdots H22^{vi} = 2.78 Å; symmetry codes: (iv) $x, y - 1, z$; (v) $-x + 1, y, -z + \frac{1}{2}$; (vi) $-x + 1, -y + 1, -z + 1$. Despite these interactions, the acetonitrile molecules show a slight disorder, as indicated by the difference electron-density map (e.g. peaks of 0.63, 0.46 and -0.52 e Å⁻³ close to acetonitrile).




Figure 2

The interaction of the acetonitrile solvent molecules with neighbouring complex molecules in (I). Dashed lines indicate the various interactions. The centroid of the C14—C19 ring is indicated by a solid dot (yellow in the electronic version of the paper). For clarity, only H atoms involved in the interactions are shown. [Symmetry codes: (iii) $-x + 1, -y + 2, -z + 1$; (iv) $x, y - 1, z$; (v) $-x + 1, y, -z + \frac{1}{2}$; (vi) $-x + 1, -y + 1, -z + 1$.]




Figure 3

C—H \cdots Cl interactions (dashed lines) in the crystal packing of (I). For clarity, only H atoms involved in the interactions are shown. [Symmetry codes: (ii) $-x + \frac{3}{2}, y + \frac{1}{2}, -z + \frac{3}{2}$; (iii) $-x + 1, -y + 2, -z + 1$; (vii) $x, y + 1, z$; (viii) $x, -y + 2, z + \frac{1}{2}$]

The shortest interactions for both Cl atoms are of the C—H \cdots Cl type (Fig. 3) [Cl1 \cdots H13ⁱⁱⁱ = 2.86 Å, Cl1 \cdots H17^{vii} = 2.75 Å, Cl2 \cdots H1ⁱⁱ = 2.60 Å, Cl2 \cdots H5^{viii} = 2.83 Å and Cl2 \cdots H27Cⁱⁱⁱ = 2.70 Å; symmetry codes: (vii) $x, y + 1, z$; (viii) $x, -y + 2, z + \frac{1}{2}$].

The Cambridge Structural Database (CSD, Version 5.34; Allen, 2002) has 18 structures containing dichlorido-[(2-(diphenylphosphanyl)benzylidene)amine]palladium(II) groups. The angle between the best planes through the benzene ring and the PdCl₂NP group range from 27.6 to 62.5°. The structure with refcode XUWHIG (Koprowski *et al.*, 2002) is almost identical to (I), only missing the *ortho*-methyl group, but crystallizes in the space group $P\bar{1}$. A fit of the Pd complex from the two structures, excluding the methyl group of (I), results in an r.m.s. deviation of 0.263 Å. As in (I), both Cl atoms attached to Pd also interact with H—C bonds. However, the interaction of the Cl atom in XUWHIG equivalent to Cl2 in (I) with the H atom in XUWHIG attached to the C atom equivalent to C25 in (I) results in the formation of centrosymmetric dimers in XUWHIG and hence in a different crystal packing compared to (I).

Experimental

All reactions were carried out under a nitrogen atmosphere using a dual vacuum/nitrogen line and standard Schlenk techniques. Solvents were dried and purified by heating under reflux and under a nitrogen atmosphere in the presence of a suitable drying agent.

Table 1
Selected geometric parameters (\AA , $^\circ$).

Pd1—Cl1	2.2770 (7)	Pd1—P1	2.2122 (8)
Pd1—Cl2	2.3664 (7)	Pd1—N1	2.049 (2)
Cl1—Pd1—Cl2	89.95 (3)	N1—Pd1—Cl1	178.15 (7)
P1—Pd1—Cl1	91.45 (3)	N1—Pd1—Cl2	91.61 (7)
P1—Pd1—Cl2	178.15 (3)	N1—Pd1—P1	87.01 (7)

For the preparation of *N*-[2-(diphenylphosphanyl)benzylidene]-2-methylaniline, 2-methylaniline (0.1030 g, 0.961 mmol) was added dropwise to a solution of 2-(diphenylphosphanyl)benzaldehyde (0.2790 g, 0.961 mmol) in dry CH_2Cl_2 (10 ml). Anhydrous magnesium sulfate (0.5 g) was added to the solution, after which the reaction was stirred at room temperature for 20 h. The resulting yellow mixture was filtered to obtain a yellow solution, which gave a yellow oil upon evaporation of the solvent (yield: 0.2990 g, 82%). IR (Nujol, ν , cm^{-1}): 1622 (C=N imine), 1593, 1585, 1505 (C=C phenyl), 1435 (P=Ph). Analysis calculated for $\text{C}_{26}\text{H}_{22}\text{NP}$: C 82.30, H 5.84, N 3.69%; found: C 82.04, H 5.93, N 3.55%.

The preparation of (I) followed the synthetic protocol of Koprowski *et al.* (2002), with minor modifications at the precipitation stage of the product. Crystals of (I) suitable for X-ray crystallographic analysis were grown by slow evaporation from an acetonitrile solution of the complex (yield: 0.0697 g, 75%). IR (Nujol, ν , cm^{-1}): 1614 (C=N imine), 1585, 1560, 1502 (C=C phenyl). Analysis calculated for $\text{C}_{26}\text{H}_{22}\text{Cl}_2\text{NPPd}$: C 56.09, H 3.98, N 2.52%; found: C 55.83, H 4.22, N 2.77%.

Crystal data



$M_r = 597.77$

Monoclinic, $C2/c$

$a = 32.350$ (3) \AA

$b = 9.9963$ (3) \AA

$c = 20.8851$ (16) \AA

$\beta = 128.423$ (13) $^\circ$

$V = 5291.3$ (12) \AA^3

$Z = 8$

Mo $K\alpha$ radiation

$\mu = 0.98$ mm $^{-1}$

$T = 100$ K

0.2 \times 0.2 \times 0.2 mm

Data collection

Agilent SuperNova diffractometer
(single source at offset, Eos
detector)

Absorption correction: multi-scan
(*CrysAlis PRO*; Agilent, 2012)
 $T_{\min} = 0.983$, $T_{\max} = 1.000$

10592 measured reflections

5391 independent reflections

4721 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.025$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.032$

$wR(F^2) = 0.075$

$S = 1.05$

5391 reflections

309 parameters

H-atom parameters constrained

$\Delta\rho_{\text{max}} = 0.75$ e \AA^{-3}

$\Delta\rho_{\text{min}} = -0.52$ e \AA^{-3}

All H atoms were placed in idealized positions and refined in riding mode, with C—H = 0.93 (aromatic) or 0.96 \AA (methyl) and $U_{\text{iso}}(\text{H}) = 1.2$ or $1.5U_{\text{eq}}(\text{C})$.

Data collection: *CrysAlis PRO* (Agilent, 2012); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *OLEX2* (Dolomanov *et al.*, 2009); software used to prepare material for publication: *OLEX2*.

The authors acknowledge the University of the Western Cape Senate Research, the South African National Research Foundation and KU Leuven International Admissions and Mobility for financial support. The authors also thank the Hercules Foundation for supporting the purchase of the diffractometer through project No. AKUL/09/0035.

Supplementary data for this paper are available from the IUCr electronic archives (Reference: SK3474). Services for accessing these data are described at the back of the journal.

References

- Agilent (2012). *CrysAlis PRO*. Agilent Technologies, Yarnton, Oxfordshire, England.
- Allen, F. H. (2002). *Acta Cryst. B* **58**, 380–388.
- Bacchi, A., Carcelli, M., Costa, M., Fochi, A., Monici, C., Pelagatti, P., Pelizzi, C., Pelizzi, G. & Roca, L. M. S. (2000). *J. Organomet. Chem.* **593**, 180–191.
- Chiririwa, H. & Muller, A. (2012). *Acta Cryst. E* **68**, m116–m117.
- Doherty, S., Knight, J. G., Scanlan, T. H., Elsegood, M. R. J. & Clegg, W. (2002). *J. Organomet. Chem.* **650**, 231–248.
- Dolomanov, O. V., Bourhis, L. J., Gildea, R. J., Howard, J. A. K. & Puschmann, H. (2009). *J. Appl. Cryst.* **42**, 339–341.
- Gao, E., Liu, C., Zhu, M., Lin, H., Wu, Q. & Liu, L. (2009). *Anticancer Agents Med. Chem.* **9**, 356–368.
- Ghilardi, C. A., Midollini, S., Moneti, S., Orlandini, A. & Scapacci, G. (1992). *J. Chem. Soc. Dalton Trans.* pp. 3371–3376.
- Koprowski, M., Sebastian, R.-M., Maraval, V., Zablocka, M., Cadierno, V., Donnadieu, B., Igau, A., Caminade, A.-M. & Majoral, J.-P. (2002). *Organometallics*, **21**, 4680–4687.
- Mogorosi, M. M., Mahamo, T., Moss, J. R., Mapolie, S. F., Slootweg, J. C., Lammertsma, K. & Smith, G. S. (2011). *J. Organomet. Chem.* **696**, 3585–3592.
- Motswainyana, W. M., Onani, M. O. & Lalancette, R. A. (2012). *Acta Cryst. E* **68**, m339.
- Newkome, G. R. (1993). *Chem. Rev.* **93**, 2067–2089.
- Nobre, S. M. & Monteiro, A. L. (2009). *J. Mol. Catal. A Chem.* **313**, 65–73.
- Ojwach, S. O., Guzei, I. A., Darkwa, J. & Mapolie, S. F. (2007). *Polyhedron*, **26**, 851–861.
- Onani, M. O., Motswainyana, W. M., Iwuoha, E. I., Darkwa, J. & Lalancette, R. A. (2010). *Acta Cryst. E* **66**, m688.
- Pellagatti, P., Carcelli, M., Costa, M., Ianelli, S., Pellizi, C. & Rogollino, D. (2005). *J. Mol. Catal. A Chem.* **226**, 107–110.
- Puri, M., Gatard, S., Smith, D. A. & Ozerov, O. V. (2011). *Organometallics*, **30**, 2472–2482.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Vaughan, T. F., Koedyk, D. J. & Spencer, J. L. (2011). *Organometallics*, **30**, 5170–5180.

supplementary materials

Acta Cryst. (2013). C69, 209–211 [doi:10.1107/S0108270113002333]

Dichlorido{N-[2-(diphenylphosphanyl)benzylidene]-2-methyl-aniline}palladium(II) acetonitrile monosolvate

William M. Motswainyana, Martin O. Onani, Jeroen Jacobs and Luc Van Meervelt

Dichlorido{N-[2-(diphenylphosphanyl)benzylidene]-2-methylaniline}palladium(II) acetonitrile monosolvate

Crystal data



$M_r = 597.77$

Monoclinic, $C2/c$

$a = 32.350$ (3) Å

$b = 9.9963$ (3) Å

$c = 20.8851$ (16) Å

$\beta = 128.423$ (13)°

$V = 5291.3$ (12) Å³

$Z = 8$

$F(000) = 2416$

$D_x = 1.501 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 5198 reflections

$\theta = 2.9\text{--}26.3$ °

$\mu = 0.98 \text{ mm}^{-1}$

$T = 100$ K

Block, yellow

0.2 × 0.2 × 0.2 mm

Data collection

Agilent SuperNova

diffractometer (single source at offset, Eos detector)

Radiation source: SuperNova (Mo) X-ray

Source

Mirror monochromator

Detector resolution: 15.9631 pixels mm⁻¹

ω scans

Absorption correction: multi-scan

(*CrysAlis PRO*; Agilent, 2012)

$T_{\min} = 0.983$, $T_{\max} = 1.000$

10592 measured reflections

5391 independent reflections

4721 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.025$

$\theta_{\max} = 26.4$ °, $\theta_{\min} = 2.9$ °

$h = -40 \rightarrow 34$

$k = -12 \rightarrow 10$

$l = -26 \rightarrow 25$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.032$

$wR(F^2) = 0.075$

$S = 1.05$

5391 reflections

309 parameters

0 restraints

Primary atom site location: structure-invariant

direct methods

Secondary atom site location: difference Fourier

map

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0255P)^2 + 11.5559P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.003$

$\Delta\rho_{\max} = 0.75 \text{ e } \text{\AA}^{-3}$

$\Delta\rho_{\min} = -0.52 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^* / U_{\text{eq}}$
Pd1	0.637890 (8)	1.06809 (2)	0.634944 (12)	0.01449 (7)
C11	0.57031 (3)	1.21650 (7)	0.58060 (4)	0.02303 (16)
C12	0.67864 (3)	1.15859 (7)	0.76645 (4)	0.02010 (15)
P1	0.60233 (3)	0.98303 (7)	0.51296 (4)	0.01688 (15)
N1	0.69700 (9)	0.9302 (2)	0.68102 (13)	0.0183 (5)
N2	0.38905 (17)	0.4779 (4)	0.1631 (3)	0.0693 (11)
C1	0.71814 (11)	0.8903 (3)	0.64898 (17)	0.0209 (6)
H1	0.7460	0.8304	0.6799	0.025*
C2	0.70380 (11)	0.9276 (3)	0.56977 (17)	0.0201 (6)
C3	0.65416 (11)	0.9765 (3)	0.50325 (16)	0.0182 (6)
C4	0.64503 (12)	1.0060 (3)	0.43027 (17)	0.0222 (6)
H4	0.6121	1.0377	0.3860	0.027*
C5	0.68440 (12)	0.9886 (3)	0.42290 (18)	0.0266 (7)
H5	0.6781	1.0107	0.3743	0.032*
C6	0.73281 (13)	0.9387 (3)	0.4874 (2)	0.0282 (7)
H6	0.7591	0.9268	0.4822	0.034*
C7	0.74248 (12)	0.9062 (3)	0.56015 (19)	0.0251 (7)
H7	0.7749	0.8700	0.6029	0.030*
C8	0.54572 (11)	1.0641 (3)	0.42268 (16)	0.0198 (6)
C9	0.54967 (12)	1.1958 (3)	0.40363 (17)	0.0243 (6)
H9	0.5818	1.2404	0.4357	0.029*
C10	0.50546 (12)	1.2584 (3)	0.33696 (18)	0.0285 (7)
H10	0.5082	1.3451	0.3237	0.034*
C11	0.45714 (12)	1.1941 (3)	0.28948 (18)	0.0302 (7)
H11	0.4276	1.2377	0.2449	0.036*
C12	0.45293 (13)	1.0648 (3)	0.30838 (18)	0.0304 (7)
H12	0.4205	1.0214	0.2765	0.036*
C13	0.49723 (11)	0.9997 (3)	0.37519 (17)	0.0250 (6)
H13	0.4943	0.9128	0.3880	0.030*
C14	0.58476 (10)	0.8086 (3)	0.50583 (16)	0.0190 (6)
C15	0.59009 (12)	0.7175 (3)	0.46123 (18)	0.0266 (7)
H15	0.6022	0.7464	0.4332	0.032*
C16	0.57752 (12)	0.5837 (3)	0.4580 (2)	0.0305 (7)
H16	0.5813	0.5231	0.4282	0.037*
C17	0.55932 (11)	0.5412 (3)	0.4995 (2)	0.0300 (7)
H17	0.5509	0.4516	0.4977	0.036*
C18	0.55362 (11)	0.6306 (3)	0.54345 (19)	0.0272 (7)

H18	0.5415	0.6011	0.5713	0.033*
C19	0.56584 (10)	0.7646 (3)	0.54659 (17)	0.0225 (6)
H19	0.5614	0.8249	0.5758	0.027*
C20	0.71923 (12)	0.8740 (3)	0.76071 (17)	0.0239 (6)
C21	0.68644 (14)	0.7899 (3)	0.7662 (2)	0.0311 (7)
H21	0.6518	0.7735	0.7203	0.037*
C22	0.70622 (14)	0.7324 (3)	0.8402 (2)	0.0330 (8)
H22	0.6857	0.6743	0.8447	0.040*
C23	0.75659 (14)	0.7626 (3)	0.9067 (2)	0.0333 (8)
H23	0.7698	0.7247	0.9568	0.040*
C24	0.78840 (13)	0.8462 (3)	0.9028 (2)	0.0347 (8)
H24	0.8224	0.8643	0.9497	0.042*
C25	0.76964 (13)	0.9056 (3)	0.82715 (19)	0.0299 (7)
C26	0.80273 (13)	0.9990 (4)	0.8220 (2)	0.0422 (9)
H26A	0.8274	1.0419	0.8740	0.063*
H26B	0.7807	1.0654	0.7811	0.063*
H26C	0.8216	0.9506	0.8077	0.063*
C27	0.43269 (18)	0.6835 (6)	0.2598 (3)	0.0827 (17)
H27A	0.4461	0.7419	0.2401	0.124*
H27B	0.4612	0.6549	0.3143	0.124*
H27C	0.4069	0.7302	0.2602	0.124*
C28	0.40812 (17)	0.5665 (5)	0.2063 (3)	0.0509 (11)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Pd1	0.01842 (11)	0.01280 (11)	0.01271 (11)	-0.00138 (8)	0.00990 (9)	0.00025 (8)
C11	0.0241 (3)	0.0242 (4)	0.0199 (3)	0.0057 (3)	0.0132 (3)	0.0033 (3)
C12	0.0244 (3)	0.0203 (3)	0.0149 (3)	-0.0019 (3)	0.0119 (3)	-0.0026 (3)
P1	0.0232 (4)	0.0133 (3)	0.0138 (3)	-0.0027 (3)	0.0113 (3)	-0.0001 (3)
N1	0.0247 (12)	0.0167 (12)	0.0152 (11)	0.0030 (10)	0.0132 (10)	0.0034 (10)
N2	0.085 (3)	0.047 (2)	0.088 (3)	-0.005 (2)	0.059 (3)	0.010 (2)
C1	0.0255 (15)	0.0170 (14)	0.0201 (14)	0.0008 (12)	0.0141 (12)	0.0020 (12)
C2	0.0300 (15)	0.0140 (13)	0.0219 (14)	0.0000 (12)	0.0188 (13)	0.0007 (12)
C3	0.0270 (14)	0.0127 (13)	0.0183 (13)	-0.0046 (12)	0.0158 (12)	-0.0032 (11)
C4	0.0319 (16)	0.0167 (14)	0.0188 (14)	-0.0067 (13)	0.0161 (13)	-0.0030 (12)
C5	0.0443 (18)	0.0191 (15)	0.0255 (15)	-0.0054 (14)	0.0262 (15)	-0.0023 (13)
C6	0.0436 (18)	0.0222 (16)	0.0367 (18)	-0.0007 (15)	0.0337 (16)	-0.0014 (14)
C7	0.0315 (16)	0.0209 (15)	0.0278 (16)	0.0005 (13)	0.0208 (14)	-0.0009 (13)
C8	0.0250 (14)	0.0198 (14)	0.0142 (13)	0.0004 (13)	0.0121 (12)	-0.0005 (12)
C9	0.0327 (16)	0.0181 (14)	0.0183 (14)	-0.0020 (13)	0.0139 (13)	-0.0010 (12)
C10	0.0425 (18)	0.0190 (15)	0.0212 (15)	0.0021 (14)	0.0184 (14)	0.0041 (13)
C11	0.0360 (17)	0.0286 (17)	0.0190 (15)	0.0066 (15)	0.0136 (14)	0.0056 (14)
C12	0.0297 (16)	0.0340 (18)	0.0190 (15)	-0.0027 (15)	0.0110 (13)	0.0000 (14)
C13	0.0292 (15)	0.0239 (16)	0.0178 (14)	-0.0033 (14)	0.0126 (13)	0.0024 (13)
C14	0.0215 (14)	0.0148 (13)	0.0149 (13)	-0.0029 (12)	0.0084 (11)	0.0012 (11)
C15	0.0365 (17)	0.0215 (16)	0.0249 (15)	-0.0097 (14)	0.0206 (14)	-0.0053 (13)
C16	0.0365 (18)	0.0190 (16)	0.0305 (17)	-0.0061 (14)	0.0182 (15)	-0.0060 (13)
C17	0.0239 (15)	0.0174 (15)	0.0356 (18)	-0.0053 (13)	0.0119 (14)	0.0033 (14)
C18	0.0222 (15)	0.0258 (16)	0.0325 (17)	-0.0045 (13)	0.0165 (14)	0.0077 (14)

C19	0.0187 (14)	0.0234 (15)	0.0237 (15)	-0.0009 (12)	0.0123 (12)	0.0024 (13)
C20	0.0359 (16)	0.0209 (15)	0.0209 (14)	0.0131 (14)	0.0207 (14)	0.0075 (13)
C21	0.057 (2)	0.0197 (15)	0.0364 (18)	0.0144 (16)	0.0385 (17)	0.0108 (14)
C22	0.047 (2)	0.0278 (17)	0.0385 (19)	0.0070 (16)	0.0337 (17)	0.0076 (15)
C23	0.050 (2)	0.0295 (18)	0.0346 (18)	0.0129 (16)	0.0332 (17)	0.0081 (15)
C24	0.0313 (17)	0.0335 (19)	0.0304 (17)	0.0083 (16)	0.0147 (15)	-0.0006 (15)
C25	0.0364 (17)	0.0268 (17)	0.0262 (16)	0.0079 (15)	0.0192 (15)	0.0049 (14)
C26	0.0329 (18)	0.052 (2)	0.0304 (18)	0.0002 (18)	0.0141 (15)	0.0054 (18)
C27	0.060 (3)	0.119 (5)	0.056 (3)	0.019 (3)	0.030 (2)	-0.028 (3)
C28	0.056 (3)	0.063 (3)	0.044 (2)	0.019 (2)	0.036 (2)	0.015 (2)

Geometric parameters (\AA , $^{\circ}$)

Pd1—Cl1	2.2770 (7)	C12—C13	1.393 (4)
Pd1—Cl2	2.3664 (7)	C13—H13	0.9300
Pd1—P1	2.2122 (8)	C14—C15	1.388 (4)
Pd1—N1	2.049 (2)	C14—C19	1.393 (4)
P1—C3	1.811 (3)	C15—H15	0.9300
P1—C8	1.809 (3)	C15—C16	1.387 (4)
P1—C14	1.811 (3)	C16—H16	0.9300
N1—C1	1.282 (3)	C16—C17	1.383 (4)
N1—C20	1.454 (3)	C17—H17	0.9300
N2—C28	1.134 (5)	C17—C18	1.373 (5)
C1—H1	0.9300	C18—H18	0.9300
C1—C2	1.462 (4)	C18—C19	1.386 (4)
C2—C3	1.405 (4)	C19—H19	0.9300
C2—C7	1.402 (4)	C20—C21	1.413 (4)
C3—C4	1.392 (4)	C20—C25	1.368 (4)
C4—H4	0.9300	C21—H21	0.9300
C4—C5	1.385 (4)	C21—C22	1.378 (4)
C5—H5	0.9300	C22—H22	0.9300
C5—C6	1.376 (4)	C22—C23	1.365 (5)
C6—H6	0.9300	C23—H23	0.9300
C6—C7	1.386 (4)	C23—C24	1.367 (5)
C7—H7	0.9300	C24—H24	0.9300
C8—C9	1.404 (4)	C24—C25	1.422 (4)
C8—C13	1.387 (4)	C25—C26	1.474 (5)
C9—H9	0.9300	C26—H26A	0.9600
C9—C10	1.379 (4)	C26—H26B	0.9600
C10—H10	0.9300	C26—H26C	0.9600
C10—C11	1.383 (4)	C27—H27A	0.9600
C11—H11	0.9300	C27—H27B	0.9600
C11—C12	1.382 (4)	C27—H27C	0.9600
C12—H12	0.9300	C27—C28	1.464 (7)
Cl1—Pd1—Cl2	89.95 (3)	C11—C12—C13	120.0 (3)
P1—Pd1—Cl1	91.45 (3)	C13—C12—H12	120.0
P1—Pd1—Cl2	178.15 (3)	C8—C13—C12	120.1 (3)
N1—Pd1—Cl1	178.15 (7)	C8—C13—H13	120.0
N1—Pd1—Cl2	91.61 (7)	C12—C13—H13	120.0

N1—Pd1—P1	87.01 (7)	C15—C14—P1	121.6 (2)
C3—P1—Pd1	106.73 (9)	C15—C14—C19	119.2 (3)
C3—P1—C14	102.78 (13)	C19—C14—P1	119.2 (2)
C8—P1—Pd1	119.56 (9)	C14—C15—H15	119.7
C8—P1—C3	107.90 (13)	C16—C15—C14	120.7 (3)
C8—P1—C14	106.41 (13)	C16—C15—H15	119.7
C14—P1—Pd1	112.13 (9)	C15—C16—H16	120.3
C1—N1—Pd1	127.53 (19)	C17—C16—C15	119.5 (3)
C1—N1—C20	115.6 (2)	C17—C16—H16	120.3
C20—N1—Pd1	116.78 (17)	C16—C17—H17	119.8
N1—C1—H1	116.0	C18—C17—C16	120.3 (3)
N1—C1—C2	127.9 (3)	C18—C17—H17	119.8
C2—C1—H1	116.0	C17—C18—H18	119.7
C3—C2—C1	124.9 (3)	C17—C18—C19	120.5 (3)
C7—C2—C1	116.1 (3)	C19—C18—H18	119.7
C7—C2—C3	119.0 (3)	C14—C19—H19	120.1
C2—C3—P1	118.1 (2)	C18—C19—C14	119.8 (3)
C4—C3—P1	122.3 (2)	C18—C19—H19	120.1
C4—C3—C2	119.4 (3)	C21—C20—N1	117.4 (3)
C3—C4—H4	119.6	C25—C20—N1	120.1 (3)
C5—C4—C3	120.7 (3)	C25—C20—C21	122.5 (3)
C5—C4—H4	119.6	C20—C21—H21	120.3
C4—C5—H5	119.9	C22—C21—C20	119.3 (3)
C6—C5—C4	120.1 (3)	C22—C21—H21	120.3
C6—C5—H5	119.9	C21—C22—H22	120.7
C5—C6—H6	119.9	C23—C22—C21	118.5 (3)
C5—C6—C7	120.1 (3)	C23—C22—H22	120.7
C7—C6—H6	119.9	C22—C23—H23	118.6
C2—C7—H7	119.7	C22—C23—C24	122.8 (3)
C6—C7—C2	120.5 (3)	C24—C23—H23	118.6
C6—C7—H7	119.7	C23—C24—H24	119.9
C9—C8—P1	120.2 (2)	C23—C24—C25	120.3 (3)
C13—C8—P1	120.0 (2)	C25—C24—H24	119.9
C13—C8—C9	119.6 (3)	C20—C25—C24	116.6 (3)
C8—C9—H9	120.3	C20—C25—C26	122.3 (3)
C10—C9—C8	119.5 (3)	C24—C25—C26	121.1 (3)
C10—C9—H9	120.3	H27A—C27—H27B	109.5
C9—C10—H10	119.5	H27A—C27—H27C	109.5
C9—C10—C11	120.9 (3)	H27B—C27—H27C	109.5
C11—C10—H10	119.5	C28—C27—H27A	109.5
C10—C11—H11	120.1	C28—C27—H27B	109.5
C12—C11—C10	119.8 (3)	C28—C27—H27C	109.5
C12—C11—H11	120.1	N2—C28—C27	178.0 (5)
C11—C12—H12	120.0		
Pd1—P1—C3—C2	45.0 (2)	C3—P1—C14—C15	-29.3 (3)
Pd1—P1—C3—C4	-141.1 (2)	C3—P1—C14—C19	150.1 (2)
Pd1—P1—C8—C9	61.7 (3)	C3—C2—C7—C6	3.0 (4)
Pd1—P1—C8—C13	-113.7 (2)	C3—C4—C5—C6	1.5 (4)

Pd1—P1—C14—C15	−143.6 (2)	C4—C5—C6—C7	−0.3 (5)
Pd1—P1—C14—C19	35.9 (2)	C5—C6—C7—C2	−2.0 (5)
Pd1—N1—C1—C2	−3.4 (4)	C7—C2—C3—P1	172.3 (2)
Pd1—N1—C20—C21	68.0 (3)	C7—C2—C3—C4	−1.8 (4)
Pd1—N1—C20—C25	−110.6 (3)	C8—P1—C3—C2	174.7 (2)
C11—Pd1—P1—C3	131.11 (10)	C8—P1—C3—C4	−11.4 (3)
C11—Pd1—P1—C8	8.46 (11)	C8—P1—C14—C15	84.0 (3)
C11—Pd1—P1—C14	−117.09 (10)	C8—P1—C14—C19	−96.6 (2)
C12—Pd1—N1—C1	−140.8 (2)	C8—C9—C10—C11	1.1 (5)
C12—Pd1—N1—C20	36.6 (2)	C9—C8—C13—C12	0.9 (4)
P1—Pd1—N1—C1	37.9 (2)	C9—C10—C11—C12	−0.6 (5)
P1—Pd1—N1—C20	−144.6 (2)	C10—C11—C12—C13	0.1 (5)
P1—C3—C4—C5	−174.3 (2)	C11—C12—C13—C8	−0.3 (5)
P1—C8—C9—C10	−176.7 (2)	C13—C8—C9—C10	−1.3 (4)
P1—C8—C13—C12	176.3 (2)	C14—P1—C3—C2	−73.1 (2)
P1—C14—C15—C16	178.6 (2)	C14—P1—C3—C4	100.7 (2)
P1—C14—C19—C18	−178.2 (2)	C14—P1—C8—C9	−170.1 (2)
N1—Pd1—P1—C3	−49.93 (11)	C14—P1—C8—C13	14.5 (3)
N1—Pd1—P1—C8	−172.58 (13)	C14—C15—C16—C17	0.2 (5)
N1—Pd1—P1—C14	61.88 (12)	C15—C14—C19—C18	1.2 (4)
N1—C1—C2—C3	−23.0 (5)	C15—C16—C17—C18	0.1 (5)
N1—C1—C2—C7	159.9 (3)	C16—C17—C18—C19	0.2 (5)
N1—C20—C21—C22	178.7 (3)	C17—C18—C19—C14	−0.9 (4)
N1—C20—C25—C24	−179.9 (3)	C19—C14—C15—C16	−0.9 (4)
N1—C20—C25—C26	1.8 (5)	C20—N1—C1—C2	179.1 (3)
C1—N1—C20—C21	−114.3 (3)	C20—C21—C22—C23	2.2 (5)
C1—N1—C20—C25	67.2 (4)	C21—C20—C25—C24	1.7 (5)
C1—C2—C3—P1	−4.7 (4)	C21—C20—C25—C26	−176.7 (3)
C1—C2—C3—C4	−178.7 (3)	C21—C22—C23—C24	−0.8 (5)
C1—C2—C7—C6	−179.8 (3)	C22—C23—C24—C25	−0.3 (5)
C2—C3—C4—C5	−0.5 (4)	C23—C24—C25—C20	−0.1 (5)
C3—P1—C8—C9	−60.4 (3)	C23—C24—C25—C26	178.2 (3)
C3—P1—C8—C13	124.2 (2)	C25—C20—C21—C22	−2.7 (5)