

HPLC Determination of Selected Flavonoid Glycosides and their Corresponding Aglycones in *Sutherlandia frutescens* Materials

Mbamalu ON^{1*}, Antunes E², Silosini N³, Samsodien H⁴ and Syce J^{1,3}

¹Discipline of Pharmacology, School of Pharmacy, University of the Western Cape, Bellville 7535, South Africa

²Department of Chemistry, University of the Western Cape, Bellville 7535, South Africa

³South African Herbal Science and Medicine Institute, University of the Western Cape, Bellville 7535, South Africa

⁴Discipline of Pharmaceutics, School of Pharmacy, University of the Western Cape, Bellville 7535, South Africa

Abstract

Sutherlandia frutescens is a popular South African plant commercially available in a range of formulations. However, reference standards for quality and stability assessment are lacking. This work reports the development and validation of a reversed phase HPLC method for the analysis of flavonoid glycosides and their corresponding aglycones in *S. frutescens* products. Five materials containing either leaf powder (LP) or spray-dried aqueous extract (SDAE) of *S. frutescens* were analysed for flavonoid content. A primary objective was to isolate non-commercially available flavonoid glycoside compounds (sutherlandins) for use as reference standards. Sutherlandins A, B, C and D were successfully isolated, and used, with other flavonoid compounds for HPLC assay development. The developed HPLC method was linear in the range of 0.2 to 60 µg/ml for quercitrin; 0.2 to 120 µg/ml for quercetin and kaempferol; 0.2 to 200 µg/ml for rutin and kaempferol-3-O-rutinoside; 4 to 180 µg/ml for sutherlandins A and D; and 4 to 200 µg/ml for sutherlandins B and C. Percentage content of sutherlandins A, B, C and D, quercetin and kaempferol in different plant materials were significantly different ($P < 0.001$). The developed HPLC method is simple, precise and robust; and can be employed for the simultaneous determination of flavonoid glycosides and aglycones for quality control of *S. frutescens* products.

Keywords: Flavonoid; HPLC assay; Sutherlandin; Quality control; Validation

Introduction

Sutherlandia frutescens L., a popular plant of the family *Fabaceae* (pea and bean family or Leguminosae), is widely used in certain parts of Southern Africa for the treatment of various ailments [1]. Randomized placebo-controlled clinical trials have been conducted to assess its safety and efficacy in HIV patients [2,3] with preparations underway for another upcoming clinical study. While a variety of formulations from this plant are commercially available and thus can be used for the clinical trial, very little is known regarding their quality and stability profiles, as well as the content of actives in such formulations.

Flavonoid glycosides have been discovered and isolated from the plant as potential actives and possible quality control markers [4,5]. Much of the biological effects of herbal medicinal remedies have been attributed to the presence of flavonoids [6,7], which are known to exist in this plant in glycosidic and aglycone forms. These different forms have differing chemistries and, as such, can be said to be representative of the different forms and conditions under which *S. frutescens* actives occur in different environments. These forms are also susceptible to extreme modification and can be readily hydrolysed, oxidized, hydroxylated, methylated, glycosylated, acylated or phenylated, giving rise to the variety of compounds within a class [8,9]. As most forms of instability in medicinal products are due to oxidative and hydrolytic reactions, advantage can be taken of flavonoid susceptibility to modification, and this can be applied in the quality control of herbal drug formulations. As such, content of flavonoid glycosides and /or corresponding aglycones can be used as an indication of product quality or stability as the flavonoid glycosides may be converted to the aglycones under unstable conditions [10]. Avula et al. reported on the quantification of flavonoid and cycloartanol glycosides in commercial preparations of *S. frutescens* [11]. However, no method has been reported for the simultaneous determination of the flavonoid glycosides and their corresponding aglycones in *S. frutescens* products.

The aim of this study therefore was to develop and validate a simple HPLC method for the simultaneous estimation of nine flavonoid compounds (two flavonoid aglycones and seven corresponding flavonoid glycosides) in *S. frutescens* formulations, with a view to employing this in resource-constrained settings for quality control and stability studies of products derived from this plant. Four of the compounds (sutherlandins A, B, C and D) are not yet commercially available and had to be isolated from the plant material for assay development and validation. Other flavonoid compounds selected for the assay were rutin, quercitrin, kaempferol-3-O-rutinoside, quercetin and kaempferol.

Materials and Methods

Materials

Samples of *S. frutescens* leaf powder (LP, batch number: BN E16794), spray-dried aqueous extracts (SDAE 1, batch number: Ferl-DST/001-1210; SDAE 2, batch number: 1674; SDAE 3, batch number: 62265 and SDAE 4, batch number: 63067) were received from suppliers within South Africa. Identification of the source plant material was by a consultant botanist of the supplying company, and a voucher specimen (PSM 602) is deposited in Parceval Laboratories, Paarl, South Africa. Commercially available reference compounds – rutin was purchased

*Corresponding author: Mbamalu ON, Discipline of Pharmacology, School of Pharmacy, University of the Western Cape, Bellville 7535, South Africa, Tel: +27219593229/2190; Fax: +278615107002; E-mail: ombamalu@uwc.ac.za

Received March 14, 2016; Accepted April 19, 2016; Published April 22, 2016

Citation: Mbamalu ON, Antunes E, Silosini N, Samsodien H, Syce J (2016) HPLC Determination of Selected Flavonoid Glycosides and their Corresponding Aglycones in *Sutherlandia frutescens* Materials. Med Aromat Plants 5: 246. doi:10.4172/2167-0412.1000246

Copyright: © 2016 Mbamalu ON, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

from Sigma Aldrich; quercetin, quercitrin, kaempferol-3-O-rutinoside and kaempferol – were purchased from Extrasynthese (Genay Cedex, France). Their corresponding glycosides were isolated from the butanol fraction of an *S. frutescens* aqueous extract. Acetonitrile (HPLC grade) and formic acid (reagent grade) were from Sigma Aldrich and Merck respectively. All water used was obtained from an in-house Milli-Q purification system (Millipore, Milford, MA).

Instrumentation and chromatographic conditions: The HPLC system for the acquisition of chromatograms and UV spectra was an Agilent 1200 series HPLC system, equipped with an in-line degassing system (G1322A, Japan), quaternary pump (G1311A, Germany), auto loading sampler (G1329A, Germany), thermostatted column compartment (G1316A, Germany) and photo diode array detector (G1315B, Germany). Chromatographic separation was obtained using a Phenomenex Luna[®] C₁₈ column (25 cm × 4.6 mm, 5 μm i.d.) with a compatible guard column, both maintained at 45°C. The mobile phase, consisting of water (0.01% formic acid) (A) and acetonitrile (0.01% formic acid) (B), was filtered through a 0.45 μm filter and degassed prior to use. The flow rate of the mobile phase was maintained at 0.8 ml/min, injection volume was 20 μL, and peaks were separated according to the following linear gradient elution: 0 to 1 min (A:B, 82%:18%); 1 to 15 min (A:B, 82%:18% to 75%:25%); 15 to 20 min (A:B, 75%:25% to 65%:35%); 20 to 25 min (A:B, 65%:35% to 40%:60%); 25 to 26 min (A:B, 40%:60% to 82%:18%); followed by an equilibration with A:B, 82%:18% from 26 to 35 minutes. The eluent was monitored at several wavelengths over a range from 250 to 380 nm, the specific wavelength of interest being 370 nm. Data acquisition and processing was carried out using the OpenLAB[™] CDS ChemStation Edition software (Agilent Technologies, Palo Alto, CA, USA).

LC-MS and LC-MS/MS analysis was conducted on a Waters Synapt G2 quadrupole time-of-flight mass spectrometer, connected to a Waters Acquity ultra-performance liquid chromatography (UPLC) and photo diode array (PDA) detector, using a method previously described by Albrecht et al. [12].

Methods

Isolation of reference compounds: Sutherlandins A to D were isolated from the *n*-butanol soluble portion of the aqueous extract of *S. frutescens* spray-dried aqueous extract (SDAE) (batch number: FerL-DST/001-1210). The crude butanol extract was subjected to reversed phase column chromatography using C₁₈ Strata[™] solid phase extraction (SPE) cartridges for the isolation of flavonoid fractions. Conditioning of the SPE cartridge was with 100% acetonitrile followed by water (0.01% formic acid). The butanol fraction (dissolved in 50% acetonitrile: water) was applied uniformly to the SPE cartridges and the column eluted using a five step gradient elution with water: acetonitrile mixtures (0.01% formic acid) of decreasing polarity (0 to 100% acetonitrile). Six major fractions were obtained; fraction 3 was found to contain the sutherlandins, and was further fractionated using HPLC. The identity of the isolated compounds was confirmed by LC-MS/MS and NMR analyses.

Preparation of calibration standards: Stock solutions of the calibration standards (rutin, quercetin, quercitrin, kaempferol, kaempferol-3-O-rutinoside, sutherlandins A to D) were prepared separately by dissolving each standard in 50% aqueous methanol solution. Dilutions were made from these stock solutions to achieve concentrations from 0.2 to 60 μg/ml for quercitrin, 0.2 to 120 μg/ml for quercetin and kaempferol, 0.2 to 200 μg/ml for rutin and kaempferol-3-O-rutinoside, 4-180 μg/ml for sutherlandins A and D, and 4 to 200 μg/

ml for sutherlandins B and C. The calibration standards were assayed in triplicate and calibration lines constructed by linear regression of plots of peak area against concentration. The prepared standards were used to validate the developed HPLC method for linearity and range, precision, limits of detection and quantitation, and ruggedness / robustness [13,14].

Sample preparation for quantification of reference compounds in plant materials: Five samples of *S. frutescens* formulations, one or more of which may be utilised in a planned randomised controlled trial, were prepared using a modification of the method previously utilized by Avula et al. [11] as follows. Weighed sample of each material: LP or SDAE material was vortexed in 2 ml of 50% aqueous methanol, and sonicated for 30 minutes. This was followed by centrifugation at 3500 rpm for 15 minutes. The procedure was repeated thrice, and respective supernatants combined. The final volume was adjusted to 8 ml with 50% aqueous methanol and mixed thoroughly. Prior to injection, appropriate dilutions were made, and the solution filtered using a 0.45 μm nylon membrane filter. The first 1.0 ml was discarded and the remaining volume collected in an LC sample vial. Fifty μl of triplicate samples was injected for each plant material.

Results and Discussion

Identification of isolated reference compounds using liquid chromatography/mass spectrometry (LC-ESI-TOF)

In this study, non-commercially available flavonoid glycoside compounds: sutherlandins A, B, C and D were isolated from *S. frutescens* SDAE, with minimum percentage content of 98.9%, 89.7%, 99.6% and 98.6%, respectively, as determined from HPLC analysis. The minimum percentage values as reported by the manufacturers for the commercially available references – quercetin, kaempferol, rutin, quercitrin and kaempferol-3-O-rutinoside – are 99.0%, 90.0%, 94.0%, 98.5% and 98%, respectively.

The identity of the isolated reference compounds was confirmed by LC-MS/MS and NMR. The MS data is shown in Figure 1. High-resolution electrospray ionisation mass spectrometry (HR ESI-MS) analysis provided the [M+H]⁺ molecular ions of the respective compounds in the positive ion mode. The protonated species [M+H]⁺, similar to those previously reported by Avula et al. [11], were therefore observed at *m/z* 741.1862, 741.1857, 725.1900 and 725.1913 amu, corresponding to the identities of sutherlandins A, B, C and D, respectively (Figure 1, Table 1).

The isolated compounds were further characterized using fragmentation patterns observed in the mass spectrum. Sutherlandins A and B showed characteristic fragments at *m/z* 609.1487 ([M+H-sugar]⁺) and 303.0484 (quercetin), while sutherlandins C and D showed characteristic fragments at *m/z* 593.1472 ([M+H-sugar]⁺) and 287.0554 (kaempferol). These ions reflect the loss of sugar from the flavonoid glycoside skeleton (Figure 1, Table 1).

Final confirmation of the identity of the isolated compounds was via NMR analyses, with results comparable to that previously reported [5].

HPLC Method

A reversed phase HPLC separation method combined with PDA detection has been developed for flavonoid analysis in *S. frutescens*. The reference compounds were separated within 30 minutes by HPLC-




Figure 1: Mass spectra of sutherlandins A to D, isolated from *S. frutescens*. Separation of the reference standards was on a Waters UPLC BEH 1.7 μm C_{18} column with dimensions 100 \times 2.1 mm, using gradient elution and flow rate of 0.35 ml/min.


Reference compound	Exact mass (amu)	[M+H] ⁺ (amu)	[M+H-sugar] ⁺ (amu)	Key fragment (amu)
Sutherlandin A	740.1800	741.1862	609.1487	303.0484
Sutherlandin B	740.1800	741.1857	609.1448	303.0498
Sutherlandin C	724.1851	725.1900	593.1472	287.0545
Sutherlandin D	724.1851	725.9130	593.1486	287.0554

Table 1: LC-ESI-MS of isolated compounds identified using positive ion mode and key fragment analysis.

DAD technique. A typical chromatogram from the assay is shown in Figure 2. The flavonoid glycosides eluted between 11 and 19 minutes, and the aglycones between 26 and 29 minutes.

Method validation

Calibration curves and linearity: The linearity for each of the



Analyte	Regression equation	Linearity range (µg/ml)	R ²	LOD (µg/ml)	LOQ (µg/ml)
Sutherlandin A	y=22.252x-9.5909	4.0 to 180.0	0.9998	1.0764	3.2617
Sutherlandin B	y=14.543x+13.149	4.0 to 200.0	0.9996	2.2984	6.9649
Sutherlandin C	y=17.890x+11.763	4.0 to 200.0	0.9990	0.3027	0.9173
Sutherlandin D	y=17.450x-1.5755	4.0 to 180.0	0.9996	0.5110	1.5484
Quercetin	y=108.85x+80.842	0.2 to 120.0	0.9970	0.1138	0.3447
Kaempferol	y=92.167x+111.36	0.2 to 120.0	0.9983	0.3309	1.0026
Rutin	y=27.929x+29.834	0.2 to 200.0	0.9989	0.8155	2.4712
Quercitrin	y=27.381x-8.4190	0.2 to 60.0	0.9997	0.1163	0.3524
K-3-O-rutinoside	y=26.799x+45.670	0.2 to 200.0	0.9991	0.0773	0.2343

Table 2: Regression equation, correlation coefficient (R²) and limits of detection (LOD) and quantitation (LOQ) for reference compounds by LC-DAD method.

reference compounds was assessed over the same concentration range as the calibration curves. The calibration curves for the nine reference standards showed a linear correlation between concentration and peak area (R²>0.99) of the detector response for the concentration ranges studied (Table 2).

Limit of detection (LOD) and limit of quantitation (LOQ): The limits of detection (LOD) and quantitation (LOQ) were calculated from the calibration curve using the equations [13,14]:

$$LOD=3.3 \times (SD/S)$$

$$LOQ=10 \times (SD/S)$$

Where, SD and S are the standard deviation of the ordinate intercept and the slope of the calibration curve, respectively. Values for the LOD and LOQ are presented in Table 2.

Stability: Samples of the reference compounds were subjected to a variety of conditions (acid/base hydrolysis, freeze/thaw cycles, ambient light and heat exposure) and injected into the HPLC system to assess the analyte in the presence of any degradants which may arise under these conditions.

Acid/base hydrolysis


The results of acid-base hydrolysis are presented in Figure 3. The

original chromatogram of each unhydrolysed compound showed only one noticeable peak (in green) in the flavonoid glycoside region for all seven flavonoid glycosides and in the aglycone region for the aglycones. On hydrolysis, the flavonoid glycosides were hydrolysed to the corresponding flavonoid aglycones (in purple), identified by comparison of their retention times and UV spectra with those of reference flavonoid aglycone compounds. This confirmed that the flavonoid glycosides are all O-glycosides, and thus prone to acid hydrolysis [15,16]. They can thus serve as references for quality control as their reactions under acid-base conditions can indicate instability.

As expected, the flavonoid aglycones were not affected by acid hydrolysis. Alkaline hydrolysis of the flavonoid glycosides led to the production of the deacylated flavonoid glycosides (navy traces, Figure 3).

Freeze-thaw cycle

The results of the freeze-thaw cycle after five consecutive freezing (at -80°C) and thawing exercises (at room temperature), are presented in Figure 4. The reference compounds showed relative stability to five successive freezing and thawing cycles, as signified by the retention times and peak sizes before and after the freeze-thaw cycle. An exception was quercetin, exhibiting instability, shown by decrease in peak size, and formation of other products. This was in contrast to




previous studies that demonstrated quercetin stability to freeze-thaw cycles, though the said studies evaluated quercetin stability in biological fluids [17,18]. Since this study utilized plant samples containing mostly flavonoid glycosides, the samples can be stored for prolonged periods in the freezer. This however may not be ideal if quercetin is to be used as a reference as it is prone to instability under freeze-thaw conditions as investigated herein, and may also be formed in cases of product instability.

Ambient light exposure

HPLC chromatograms of samples subjected to sunlight exposure for one month (Figure 5) indicates that the flavonoid glycosides (sutherlandins A, B, C and D) showed relatively good stability with respect to the retention times (i.e., still eluted at previous retention times) while the aglycones were not stable, with additional peaks, possibly from degraded products, arising in the chromatograms for

quercetin and kaempferol (Figure 5). Retention times for the flavonoid glycosides were not affected by exposure to ambient light; however, the peak response for all the seven flavonoid glycosides decreased to between 45 and 97% of the original.

The peaks for the flavonoid aglycones, quercetin and kaempferol, were however not detected in the chromatograms following light exposure, perhaps due to the photo-oxidation of these compounds [19]. The stability of flavonoids to light is known to depend on the nature of the hydroxyl group attached to C₃ of the flavonoid structure, where the absence or glycosylation of the hydroxyl group confers some degree of stability not seen in flavonoid molecules with a hydroxyl group at C₃ [20] as is the case with quercetin and kaempferol. The seven flavonoid glycosides under study are all glycosylated at position C₃, conferring on them some degree of photostability not obtained with the non-glycosylated flavonoid aglycones, quercetin and kaempferol. This may



explain why the flavonoid glycosides, though affected by ambient light exposure which served to reduce the peak response, were affected to a lesser degree than the flavonoid aglycones of lower photostability due to the presence of a free hydroxyl group at C₃. The 'lower-stability' flavonoid aglycones underwent degradation to such an extent that their peaks were not detected in the chromatograms obtained following ambient light exposure. The lower photostability of the flavonoid aglycones is ascribed to a greater triplet state population *cum* higher singlet oxygen reactivity [20].

All the compounds exhibited instability on exposure to light; ideally, samples containing these compounds should not be exposed to ambient light conditions as tested here. This also shows that the flavonoid compounds are susceptible to light and can therefore be used as reference compounds to monitor quality and stability of *S. frutescens* products exposed to light.

Heat exposure

Following heat exposure (for 3 hours at 60 degrees Celsius), all the flavonoid glycosides and aglycones were still detected at their respective retention times (Figure 6) indicating that heat did not cause exhaustive degradation of the compounds.

The peak response was however reduced to between 4 and 84% of the original for the flavonoid glycosides and aglycones under study. This is in line with a study by Manach et al. [21] which reported a 60%

loss of quercetin in apple juice stored for 9 months at 25°Celsius. The flavonoid glycoside, kaempferol-3-O-rutinoside, where peak response increased to 106.8% after heat exposure was an exception from the other flavonoids in the group. However, such increase may also be viewed as being due to some form of instability or an indication of compromised quality. Ideally, samples containing these compounds should not be exposed to heat conditions as tested here. The flavonoid compounds, having shown susceptibility to heat, can therefore be used as markers for quality and stability assessment of *S. frutescens* products exposed to heat.

Precision and accuracy

The precision of the analytical method was determined by assaying 6 samples all at a concentration of 60 µg/ml; the highest concentration for the analyte with the lowest upper limit. Intra-day and inter-day precision were assessed by assay of six replicate samples on the same day, over three different days. The relative standard deviations (% RSD) were calculated for the standard samples at the concentrations assayed and are presented in Table 3.

The results of intraday and interday assays were consistent, with % RSD less than 1% for all reference compounds (Table 3). The precision and accuracy data are therefore within acceptable limits for *S. frutescens* assay.




Figure 5: HPLC chromatograms (at 370 nm and 250 nm) for reference compounds after sunlight exposure for one month. SA: sutherlandin A; SB: sutherlandin B; SC: sutherlandin C; SD: sutherlandin D; K: kaempferol; Ko: kaempferol-3-O-rutinoside; Q: quercetin; R: rutin; Qr: quercitrin.

Reference	Ave peak area	% RSD	Intraday Ave	Intraday % RSD	Interday Ave	Interday % RSD
Sutherlandin A	1325.99	0.998	1341.60	0.061	1328.03	0.442
Sutherlandin B	916.62	0.442	933.47	0.469	909.18	0.818
Sutherlandin C	1167.74	0.111	1130.98	0.452	1136.91	0.629
Sutherlandin D	1085.46	0.205	1087.62	0.056	1087.89	0.959
Quercetin	7134.14	0.146	7063.00	0.433	6987.52	0.327
Kaempferol	5952.61	0.242	5967.14	0.090	5989.19	0.494
Rutin	1836.46	0.738	1861.07	0.161	1853.79	0.355
Quercitrin	1646.19	0.045	1659.16	0.048	1662.54	0.457
Kaempferol-3-o-rutinoside	1759.65	0.019	1761.84	0.204	1782.92	0.502

Table 3: Intra- and inter-day precision analyses for reference compounds at 370 nm.

Robustness and ruggedness

Evaluation of method robustness was considered in mobile phase development. The robustness and ruggedness of the method were investigated by varying chromatographic parameters such as the flow rate and temperature.

A decrease in the flow rate of the mobile phase (from 0.8 to 0.5 ml/minute) resulted in elongation of the retention times for all the reference compounds (Figure 7a), while changing the temperature of the column compartment from 40°C to room temperature did not cause significant changes in the peak sizes and retention times, showing that the developed method is robust to temperature changes (Figure 7b).

System suitability testing

System suitability parameters were calculated from the chromatograms obtained for each of the reference compounds during the study (Table 4). Values of the capacity factor (k') ranged from 30 to 71, resolution (R_s) from 49 to 245 and tailing factor (T) from 1.0574 to 1.2563. The suitability of the chromatographic system in terms of column efficiency, resolution and precision, for flavonoid analysis in *S. frutescens* is therefore assured.

Quantification of reference compounds in plant materials

The aim of this study was to develop a simple HPLC assay for the simultaneous determination of flavonoid glycosides and their corresponding aglycones in *S. frutescens* formulations.




Figure 6: HPLC chromatograms (at 370 nm) for reference compounds after heat exposure for 3 hours at 60°C. SA: sutherlandin A; SB: sutherlandin B; SC: sutherlandin C; SD: sutherlandin D; K: kaempferol; Ko: kaempferol-3-O-rutinoside; Q: quercetin; R: rutin; Qr: quercitrin.




Figure 7a: HPLC chromatograms at 370 nm for a mixture of the nine reference compounds, demonstrating the effect of flow rate: a) 0.8 ml/min and b) 0.5 ml/min.




Figure 7b: HPLC chromatograms at 370 nm for a mixture of the nine reference compounds, with column temperature at: a) 45°C and b) room temperature.

Compound	k'	R _s	T
Sutherlandin A (SA)	31.5870	81.5940	1.0714
Sutherlandin B (SB)	34.4686	83.9060	1.0574
Sutherlandin C (SC)	40.6013	61.4830	1.1421
Sutherlandin D (SD)	42.6056	48.9990	1.0843
Quercetin (Q)	66.3686	244.5770	1.1981
Kaempferol (K)	71.2636	-	1.2563
Rutin (R)	29.7370	85.0840	1.1117
Kaempferol-3-O-rutinoside (Ko)	38.3600	97.0670	1.1065
Quercitrin (Qr)	45.3070	97.9100	1.0903

Table 4: HPLC system suitability parameters for the reference compounds.




Figure 8: HPLC chromatograms of the five *S. frutescens* materials in 50% aqueous methanol solution. The numbers 1 to 6 represent sutherlandin A, sutherlandin B, sutherlandin C, sutherlandin D, quercetin and kaempferol, respectively.

Five *S. frutescens* materials (leaf powder, LP; and four spray-dried aqueous extracts, SDAE : SDAE 1, SDAE 2, SDAE 3 and SDAE 4) were analysed and their flavonoid glycoside and aglycone contents determined. The HPLC separation of reference compounds in the five *S. frutescens* materials analysed is presented in Figure 8. The flavonoid glycosides, sutherlandins A, B, C and D, represented by peaks 1, 2, 3 and 4, respectively, were detected in all the samples assayed. The flavonoid aglycones, quercetin and kaempferol, represented by peaks 5 and 6, respectively, were only detected in two of the SDAE materials and not in the LP material. Three of the nine compounds, rutin, quercitrin and kaempferol-3-O-rutinoside were not conclusively identified in the plant materials assayed. While peaks with similar retention times and UV-spectra as some of these three compounds were detected in some of the *S. frutescens* materials, materials, they were not isolated and so their identities could not be confirmed by further analytical techniques.

The average peak area was used to quantify the reference compounds contained in each of the five *S. frutescens* materials. Values of the flavonoid content in the five *S. frutescens* materials, expressed as percentage content (w/w), and are shown in Table 5. The results show

Reference compound	LP	SDAE 1	SDAE 2	SDAE 3	SDAE 4
SA	0.657 ± 0.006	0.724 ± 0.013	0.250 ± 0.003	0.808 ± 0.027	0.090 ± 0.002
SB	0.562 ± 0.042	0.746 ± 0.008	0.949 ± 0.006	1.255 ± 0.022	0.700 ± 0.002
SC	1.970 ± 0.011	1.506 ± 0.003	1.355 ± 0.001	1.678 ± 0.040	1.387 ± 0.017
SD	0.642 ± 0.022	0.553 ± 0.001	0.876 ± 0.023	0.972 ± 0.045	0.828 ± 0.025
Q	ND	ND	0.030 ± 0.000	< LOQ	0.018 ± 0.000
K	ND	ND	0.263 ± 0.000	< LOQ	0.068 ± 0.000

Each value is expressed as average percentage content (mg flavonoid x minimum purity of flavonoid reference/mg plant material) ± SD (n=3). Means within a row for each flavonoid compound are significantly different (p<0.001). Key: SA: sutherlandin A; SB: sutherlandin B; SC: sutherlandin C; sutherlandin D; Q: quercetin; K: kaempferol; ND: Not detected; <LOQ: less than LOQ.

Table 5: Percentage (w/w) of reference standards found in the different *S. frutescens* extracts. Results are presented as percentage content (w/w).

that of the two possible forms in which the flavonoids can exist in plant materials (glycoside and aglycone), the flavonoid glycosides were more abundant than the flavonoid aglycones in each of the assayed materials.

The flavonoid aglycone content (0.02 to 0.26%) was found to be lower than that of the flavonoid glycosides (0.09 to 1.97%), and in some cases, below the quantification limit. The spray-dried extracts contained 0.02 to 0.26% of the flavonoid aglycones, detected in half of the SDAE investigated while these compounds were not detected in the LP samples. This suggests that the flavonoid aglycone content may be increased by processing parameters, such as is used in the preparation of the spray-dried extracts. However, no collaboration of this was found in existing literature.

In all the plant materials assayed, sutherlandin C was found to have the highest abundance, and so can be assayed for in all the *S. frutescens* materials. The least abundant flavonoid glycoside was not the same in all the plant samples assayed, with different materials showing non-uniformity in the flavonoid of least abundance. For instance, in the LP and SDAE 1 materials, the flavonoid glycosides of least abundance were sutherlandin B and sutherlandin D, respectively, while in the other three spray-dried materials, the flavonoid glycoside of least abundance was sutherlandin A.

Three out of the five materials did not contain quantifiable levels of the aglycones, quercetin and kaempferol. In the two spray-dried materials that contained the aglycones (SDAE 2 and SDAE 4), kaempferol content was more than quercetin content. The sum of the analysed flavonoids in the plant materials exceeds that obtained in other studies of total flavonoids [22-24], and suggests that the flavonoid content in the *S. frutescens* materials is sufficient for flavonoid use as reference marker compounds. Overall, the levels of the flavonoid glycosides and aglycones in the different *S. frutescens* materials assessed varied significantly (p<0.001). The assessment of flavonoid glycoside (sutherlandin A, B, C and D) content can therefore be used to differentiate between *S. frutescens* products.

The levels of the flavonoid aglycones (quercetin and kaempferol) in the analysed samples were quite low and in some cases, absent, thus making quantification of these compounds a challenge. The ratio of the flavonoid glycosides to the total flavonoids (glycoside + aglycone) was at least 90% in the samples studied. Therefore, the use of the flavonoid aglycones (and not the glycosides) as markers for quality control and stability assessment of *S. frutescens* LP or SDAE materials may not be justifiable or feasible using the specific assay employed in the present study. This however, can be expected in samples extracted using more polar solvents like methanol and water. It may well be that extraction with less polar solvents would yield more of the aglycones than the glycosides.

Conclusions

The simple, validated HPLC-DAD method reported was found to be appropriate for separation and quantification of flavonoid glycosides and their corresponding aglycones in different formulations of *S. frutescens*. The LP and SDAE materials all contained all four marker compounds, i.e., sutherlandins A, B, C and D, with sutherlandin C being the most abundant in all the five materials. The two flavonoid aglycones, quercetin and kaempferol, were detected in two of the SDAE materials, and not in the LP material. There were more flavonoid glycosides than aglycones in all the plant materials, suggesting that the former may be more suitable as marker compounds. The use of the flavonoid glycosides as markers is therefore recommended for assessment of *S. frutescens* materials.

The aglycones can also be used as marker compounds because their presence, as products of flavonoid glycoside breakdown, can indicate instability of the plant materials. However, use of the flavonoid aglycones levels alone, without the flavonoid glycosides levels as well, is not recommended. Assay of all six flavonoids employed here is preferred to the assay of a single flavonoid as it gives the profiles and levels of the flavonoids in the different materials. This in turn can indicate quality status of different *S. frutescens* products.

Acknowledgements

The authors acknowledge the Department of Science and Technology, South Africa, grant number DST/ICON 167/2010 dated 4 November 2010 and revised on 10 Oct 2013 for funding assistance. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests

The authors declare no competing interests.

References

- Ojewole JAO (2004) Analgesic, anti-inflammatory and hypoglycemic effects of *Sutherlandia frutescens* R. Br. (variety Incana E.MEY.) [Fabaceae] shoot aqueous extract. *Methods and Findings in Experimental and Clinical Pharmacology* 26: 409-416.
- Johnson Q, Syce J, Nell H, Rudeen K, Folk WR (2007) A randomized, double-blind, placebo-controlled trial of *Lessertia frutescens* in healthy adults. *PLoS Clin Trials* 2: e16.
- Wilson D, Goggin K, Williams K, Gerkovich MM, Gqaleni N, et al. (2015) Consumption of *Sutherlandia frutescens* by HIV-seropositive South African adults: an adaptive double-blind randomized placebo controlled trial. *Plos One* 10: 0128522.
- Fu X, Li XC, Smillie TJ, Carvalho P, Mabusela W, et al. (2008) Cycloartane glycosides from *Sutherlandia frutescens*. *J Nat Prod* 71: 1749-1753.
- Fu X, Li XC, Wang YH, Avula B, Smillie TJ, et al. (2010) Flavonol glycosides from the south African medicinal plant *Sutherlandia frutescens*. *Planta Med* 76: 178-181.
- Cai Y, Luo Q, Sun M, Corke H (2004) Antioxidant activity and phenolic compounds of 112 traditional Chinese medicinal plants associated with anticancer. *Life Sci* 74: 2157-2184.
- Atawodi SE (2005) Antioxidant potential of African medicinal plants. *African Journal of Biotechnology* 4: 128-133.
- Schwinn KE, Davies KM (2004) Flavonoids In: Davies KM (Ed.), *Plant pigments and their manipulation*. Blackwell, pp: 92-149.
- Winkel-Shirley BSJ (2006) The biosynthesis of flavonoids In: Grotewold E (Ed.), *The Science of Flavonoids*. Springer, pp: 71-95.

Citation: Mbamalu ON, Antunes E, Silosini N, Samsodien H, Syce J (2016) HPLC Determination of Selected Flavonoid Glycosides and their Corresponding Aglycones in *Sutherlandia frutescens* Materials. *Med Aromat Plants* 5: 246. doi:10.4172/2167-0412.1000246

- Deng S, West BJ, Jensen CJ (2011) Thermal degradation of flavonoid glycosides in noni leaves during roasting. *Advanced Journal of Food Science and Technology* 3: 155-159.
- Avula B, Wang YH, Smillie TJ, Fu X, Li XC, et al. (2010) Quantitative determination of flavonoids and cycloartanol glycosides from aerial parts of *Sutherlandia frutescens* (L.) R. BR. by using LC-UV/ELSD methods and confirmation by using LC-MS method. *J Pharm Biomed Anal* 52: 173-180.
- Albrecht CF, Stander MA, Grobbelar MC, Colling J, Kossman J, et al. (2012) LC-MS-based metabolomics assists with quality assessment and traceability of wild and cultivated plants of *Sutherlandia frutescens* (Fabaceae). *South African Journal of Botany* 82: 33-45.
- ICH (1997) International Conference on Harmonization of technical requirements for registration of pharmaceuticals for human use. Section Q2B: Validation of analytical procedures: methodology.
- Shahir GA (2003) Validation of high-performance liquid chromatography methods for pharmaceutical analysis. Understanding the differences and similarities between validation requirements of the US Food and Drug Administration, the US Pharmacopeia and the International Conference on Harmonization. *J Chromatogr A* 987: 57-66.
- Bohm BA (1998) *Introduction to flavonoids*. Harwood Academic Publishers, The Netherlands.
- Sun Y, Zhang X, Xue X, Zhang Y, Xiao H, et al. (2009) Rapid identification of polyphenol C-glycosides from *Swertia franchetiana* by HPLC-ESI-MS-MS. *J Chromatogr Sci* 47: 190-196.
- Li N, Liu C, Mi S, Wang N, Zheng X, et al. (2012) Simultaneous determination of oleanolic acid, *p*-coumaric acid, ferulic acid, kaempferol and quercetin in rat plasma by LC-MS-MS and application to a pharmacokinetic study of *Oldenlandia diffusa* extract in rats. *Journal of Chromatographic Science*, pp: 1-8.
- Duppala V, Dash RP, Jivrajani MN, Thakur SK, Ravat NM, et al. (2013) Simultaneous quantification of glibenclamide, simvastatin, and quercetin by using LC-UV method and its application to pharmacokinetic study in rats. *Chromatography Research International*, Article ID 768160.
- Dall'Acqua S, Miolo G, Innocenti G, Caffieri S (2012) The photodegradation of quercetin: relation to oxidation. *Molecules* 17: 8898-8907.
- Smith GJ, Thomsen SJ, Markham KR, Andart C, Cardon D (2000) The photostabilities of naturally occurring 5-hydroxyflavones, flavonols, their glycosides and their aluminium complexes. *Journal of Photochemistry and Photobiology A: Chemistry* 136: 87-91.
- Manach C, Scalbert A, Morand C, Rémésy C, Jiménez L (2004) Polyphenols: food sources and bioavailability. *American Journal of Clinical Nutrition* 79: 727-747.
- Kaur S, Mondal P (2014) Study of total phenolic and flavonoid content, antioxidant activity and antimicrobial properties of medicinal plants. *Journal of Microbiology and Experimentation*, 1: 00005.
- Srivastava MP, Tiwari R, Sharma N (2013) Assessment of phenol and flavonoid content in the plant materials. *Journal of New Biological Reports* 2: 163-166.
- Samatha T, Shyamsundarachary R, Srinivas P, Swamy NR (2012) Quantification of total phenolic and total flavonoid contents in extracts of *Oroxylum indicum* L. Kurz. *Asian Journal of Pharmaceutical and Clinical Research* 5: 177-179.

OMICS International: Publication Benefits & Features

Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

Special features:

- 700 Open Access Journals
- 50,000 Editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus, Google Scholar etc.
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submit>